# Academic Achievement Reflected the Quality of Life and Neurocognitive Status in Malaysian Primary School Children

Gisely Vionalita<sup>1</sup>, I. Zalina<sup>2</sup> and W. A. Asim<sup>2</sup>

<sup>1</sup>Department of Public Health, Faculty of Health Sciences, Universitas Esa Unggul, Jakarta <sup>2</sup>BRAINetwork Centre for Neurocognitive Sciences, School of Health Sciences, Universiti Sains Malaysia, Kubang Kerian Kelantan 16150

Keywords: CANTAB, primary school children, cognitive status, academic achievement.

Academic achievement is not the only indicator of achievement in school. The neurocognitive profile, as Abstract: measured by the quality of life (QoL) and neurocognitive status are important markers that make academic achievement more relevant about optimum brain utilization. The main objective of this study was to determine the relationship between quality of life, neurocognitive status and academic achievement in a sample of children in a Malaysian primary school. QoL was measured using three versions of the TNO-AZL quality of life (TACQOL) questionnaire from the children's (CV), parents' (PV) and teachers' (TV) perspectives. Neurocognitive status was assessed in two domains, executive function and visual memory using Cambridge Neuropsychology Tests Automated Battery (CANTAB®). A convenience sampling was undertaken for all 95 Standard One Zainab 2 Primary School children (7 Years Old), 95 parents and 4 teachers. Regarding executive function, 42.1% of children experienced difficulties in an extra dimensional shift which measures attention flexibility in accepting a new rule. However, they performed well in theintraextra dimensional shift, indicating that most children responded positively to experiential learning. For visual memory, 49.5% of children experienced some difficulties regarding their short term visual memory. The relationship between QoL, neurocognitive status and academic achievement of the children showed that only 'cognitive complaint' and 'negative mood' had a significant linear positive relationship with academic achievement. Thus, this study has shown that academic achievement does not necessarily reflect the neurocognitive status of children implying that some neurocognitive problems remain undetected. A more meaningful assessment of academic achievement, in primary schools, should include an assessment of both QoL and neurocognitive profile.

## **1** INTRODUCTION

Current research indicates that contrary to previous concepts which postulate that learning occurs at specific periods of the developing brain, there is evidence that both the developing and the mature brain are structurally altered when learning occurs. These structural changes are believed to encode the learning process in the brain. Studies have found that direct contact with a stimulating physical environment and an interactive social group can alter the structure of nerve cells and of the tissues that support them (Breslow, 1999). The nerve cells develop a greater number of synapses through which they communicate with each other, and the structure of the nerve cells themselves is correspondingly altered.

Quality of life (QoL) is a multidimensional construct referring to the subjective perception of physical, mental, social, psychological and functional aspects of well-being and health (Cella and Tulsky, 1990; Felce and Perry, 1995; Hofstede, 1984). From an epidemiological point of view, information on the quality of life is important to define health problems and to evaluate the wellbeing and functioning of populations (Bullinger et al., 2008). Indeed, QoL conceptualised as the individual's self-evaluation of their health status is an important criterion in evaluating health and health care. Until now, few systematic attempts have been made to develop instruments to assess the QoL of children using such a conceptualisation (Vogels et

Vionalita, G., Zalina, I. and Asim, W.

- Academic Achievement Reflected the Quality of Life and Neurocognitive Status in Malaysian Primary School Children. DOI: 10.5220/0009950126632669
- In Proceedings of the 1st International Conference on Recent Innovations (ICRI 2018), pages 2663-2669 ISBN: 978-989-758-458-9
- Copyright © 2020 by SCITEPRESS Science and Technology Publications, Lda. All rights reserved

*al.*, 2000). In this study the QoL is one of the variables that will be used in the self-evaluation of wellness and the tool chosen is the Netherlands Organisation for Applied Scientific Research Academic Medical Centre (TNO-AZL) child quality of life (TACQOL) (Vogels et al., 1998; Kamphuis et al., 1997; Kolsteren et al., 2001; Vogels et al., 1999; Bernard et al., 2009) questionnaire. In this case, the QoL, as assessed by the TACQOL, is defined as children's health status, weighted by the emotional response of their health status problems from their perspectives.

Indeed, children are the best source of concerning their feelings information and evaluations even though they may be lacking in their vocabulary and reading skills. The TACQOL considers this and has been designed with simple language and can be completed in a short period. TACQOL is a multidimensional instrument, with 7 subscales. The subscales covered by the TACOOL are based on a review of the literature, discussions with experts (child psychologists, paediatricians) and statistical testing (Vogels et al., 2000). The TACQOL was constructed to enable a systematic, valid and reliable description of the quality of life of children aged 6 till 15 by the children themselves or their parents (Arnould et al., 2004).

In this study, the neurocognitive profiling and the assessment of the neurocognitive status of primary school children will focus on the executive functions of the prefrontal cortex in relation to the limbic system and how they impact on the academic achievement and QoL of these children. The neurocognitive evaluation of school children was carried out using the Cambridge Neuropsychological Tests Automated Battery (CANTAB<sup>®</sup>) (Luciana, 2003; Cognition, 2007; Sharma, 2013; Robbins et al., 1994; Sahakian and Owen, 1992; De Luca et al., 2003).These tests are comparable across cultures as they have the advantage of being languageindependent and culture-free.

In this study, the CANTAB<sup>®</sup> was chosen for the assessment of neurocognitive status as it allows for a complete assessment of the neurocognitive profile through an extensively validated neuropsychological battery of tests. It is highly portable and uses a touch screen computer system which can be easily applied to all respondents. The CANTAB<sup>®</sup> consists of several well-validated tests, each of which taps one or more cognitive domains, and produces some outcome variables. There are six domains which are assessed in CANTAB<sup>®</sup>. In this study, two tests from CANTAB<sup>®</sup>were used from two domains (executive function and visual memory) which are closely

related to learning and memory. IED Intra-Extra Dimensional Set Shift (IED) (Lukowski et al., 2010; Lythe et al., 2005; Collinson et al., 2014) was used to assess executive function of the children while Paired Associates Learning (PAL) (de Rover et al., 2011; O'Connell et al., 2004; Barnett et al., 2015) was used to assess the visual memory of the children.

IED is primarily sensitive to changes affecting the frontostriatal areas of the brain. It is sensitive to cognitive changes associated with visual discrimination and attentional set formation maintenance, shifting and flexibility of attention (Heinzel *et al.*, 2010).

The PAL (Paired Associates Learning) was chosen from the CANTAB<sup>®</sup> as one of two tests that were carried out in this study. PAL measures visual memory, and it assesses episodic memory, new learning and is sensitive to medial temporal lobe functioning. Thus, in this study, PAL allowed for the assessment of the learning of items (as syllables, digits or words) in pairs so that one member of the pair evokes recall of the other.

In the context of this study, academic achievement is defined as the outcome of education in the form of knowledge, skills and attitude, which a student has achieved through attaining specific educational goals. Examinations or continuous assessment commonly measure academic achievementbut there is no general agreement on how it is best tested or which aspects are most important. It is one of the indicators that can describe the school-based performance of children.

# 2 RESEARCH METHOD

This cross-sectional study involved the observation of all members of a population, at one specific point in time. In this case, the study involved all Standard One children (7 Years Old) of Zainab 2 Primary School, Kota Bharu, Kelantan and included their parents as well as teachers.

## 2.1 Quality of Life

Quality of life was assessed from primary data obtained from a translated and validated TACQOL questionnaire with three versions. A parent and teacher version of the TACQOL questionnaire was constructed for parents and teachers to assess the quality of life of the children from their perspectives. The TACQOL - Child Version (TACQOL-CV) was constructed for children to assess themselves. The TACQOL – Parent/Teacher Version (TACQOL-P/TV) specifically asked parents and teachers to evaluate and assess the children's feelings about their physical, emotional and cognitive status. The major difference between the versions (child, parent and teacher) was regarding the perspective of the questions asked. The parent and teacher questionnaires used the following format: "has your child had.... or did your child have...?", while the child questionnaire used the format: "have you had .... or did you have...?".

The English version of the TACQOL was forward translated into Bahasa Malaysia and subsequently back-translated into English. This questionnaire had 56 items, with a Likert scale type of assessment (never, occasionally, only within the past weeks, often and always). The TACOOL questionnaire covered seven subscales: physical complaints and motor functioning (physical), autonomous functioning (daily living), social functioning (social), cognitive functioning and positive moods and negative moods (Verrips et al., 1999). All respondents were interviewed and observed within the school setting using the translated and validated questionnaire. Informed consent was obtained from all respondents of the study, following which the questionnaire was completed by 95 Standard One children of Zainab 2 Primary School, Kota Bharu Kelantan, 95 parents of the same children and 4 teachers who taught these Standard One children.

### 2.2 Neurocognitive Status

Neurocognitive function was analysed from primary data obtained from two tests: IED (Intra Extra Dimensional Set Shift) and PAL (Paired Associates Learning) from within the CANTAB®battery tests. These two tests were sequentially administered to the respondents by trained data collectors. The tests were administered to 95 children in a conducive and quiet place with minimal distractions. These tests were completed within 20 minutes and were simple to understand. The same instructor for both tests supervised all the children.

The IED screened for executive function in nine stages. Stage 1 started with two simple, colour-filled shapes and required children to learn and touch the correct stimuli to show simple discrimination learning. In Stage 2, the contingencies were reversed to show reversal learning. In Stage 3, the second dimension was introduced, with shapes and lines together as a distraction. The contingencies were not changed in Stage 4 but included overlapping and simple discrimination. Reversal learning occurs again in Stage 5. New compound stimuli were presented in Stage 6. Subjects were required to be attentive to the previously relevant dimension of shapes and learn which of the two exemplars were correct. Stage 7 completed an interdimensional shift. In Stage 8, children were required to shift attention to the previously irrelevant dimension and learn which of the two exemplars in this dimension were now correct. Thisgave a good indication of attentional flexibility. In Stage 9, the contingencies were again reversed.

PAL screened the visual memory ability in 5 separate stages. The children were required to remember the shapes displayed on the screen. The test started with 1 shape and then progressed to 2 shapes, 3 shapes, 6 shapes and 8 shapes. For each stage, boxes were displayed on the screen. All were opened in a randomized order. Two or more of them contained a pattern.

The patterns in the boxes were then displayed in the middle of the screen, one at a time and the subject was asked to touch the box where the pattern was originally located.Each stage had up to ten trials. If the subject made an error, the patterns were presented to remind the subject of their locations. When the subjects correctly identified allthe locations, they proceeded to the next stage. If the subject was unable to complete a stage correctly, the test was terminated.

The CANTAB<sup>®</sup>battery of tests wascomputerized and provided automatic results that were presented based on the requested outcome measurement. From the summary template definition, the researcher was able to select the outcome measurement that was required. The results were automatically calculated within the datasheet format for further statistical analysis. In this study, the outcome measurement for executive function and visual memory were a measure of the total errors.

### 2.3 Academic Achievement

Academic achievement was analyzed from secondary data obtained from the school database of the Zainab 2 Primary School as shown in Figure 1. The data was obtained from the average of mid-year examination of 2011. The data provided information on six subjects taught within the Standard One school curriculum.



#### 3 RESULTS

The average scores of the subscales in quality of life assessments from children's, parents' and teachers' perspectives together with a total error of both the IED and PAL tests were assessedregarding their relationship to the average academic achievement of the Standard One children from Zainab 2 Primary School. Simple linear regressions followed by multiple linear regressionwere appliedtoanalyzing the relationship between QoL, neurocognitive status and academic achievement. As a requirement for applying the multiple linear regression, simple linear regression was applied first to check the significance of the factors. From the simple linear regression for univariable analysis, it was found that all the variables were significant factorsfor academic achievement (p<0.25) (Table 1). Following this, all the variables were subjected to multiple linear regression for variable selection. It was seen that only 'cognitive' and 'negative mood' had a

significant linear relationship with academic achievement (p<0.05) (Table2).

Multiple linear regression analysis showed a significant linear positive relationship between both cognitive and negative domains of the quality of life assessment with the academic achievement. One single improvement in cognitive problem increased the academic achievement by a score of 0.78 and for one single improvement in negative problem the academic achievement improved by 1.27. However, there was no significant linear relationship between neurocognitive status and academic achievement, indicating that the academic achievement does not fully represent the neurocognitive status of the children.

Table 1: Sin	nple linear reg	ression for	variables
--------------	-----------------	-------------	-----------

Variable	b <sup>a</sup> (95% CI)	P-
		Value
Physical	0.78 (-0.22, 1.79)	0.126
Motor	0.68 (-0.40, 1.75)	0.215
Autonomy	1.01 (0.08, 1.94)	0.003
Cognitive	1.43 (0.85, 2.02)	0.000
Social	0.99 (-0.15, 2.13)	0.087
Positive Mood	0.77 (-0.15, 1.68)	0.098
Negative Mood	1.91 (1.18, 2.64)	0.000
Total Error IED	-0.16 (-0.36, 0.26)	0.090
Total Error PAL	-0.26 (-0.53, 0.02)	0.052

*P-value significant at <0.25* 

Table 2: Relationship between variables with academic achievement in Standard One children of Zainab 2 Primary School

b <sup>a</sup> (95% CI)	t-stat	P-value
0.78 (0.03,	2.08	0.041
1.53)	2.68	0.009
1.27 (0.33,		
2.22)		
	0.78 (0.03, 1.53) 1.27 (0.33,	0.78 (0.03, 2.08 1.53) 2.68 1.27 (0.33,

<sup>a</sup>adjusted regression coefficient <sup>b</sup>P-value significant at <0.05

Forward multiple linear regression applied. Model assumptions are fulfilled. There were no interactions amongst independent variables. No multicollinearity detected.

Coefficient of determination (R2) = 0.254

### Final model equation

Academic achievement= 31.26 + 1.27 \* Negative + 0.78 \* Cognitive

## **4 DISCUSSION**

The objective of this study was to evaluate the relationship between quality of life, neurocognitive status and academic achievement. As mentioned earlier, the average scores of the subscales in quality of life assessments from children's, parents' and teachers' perspectives together with total error of both the CANTAB<sup>®</sup> IED and CANTAB<sup>®</sup> PAL tests were assessed in terms of their relationship to the academic achievement of the Standard One Zainab 2 Primary School children.

From the statistical analysis, involving seven in OoL and two domains in subscales neurocognitive status, it was found that a significant relationship existed between two subscales of the QOL. Five other subscales were not significantly related to academic achievement. Those five subscales are physical, motor, autonomy, social and positive moods. These findings correlate well with De Oliveira Filho and Vieira (2007), where the researchers were also unable to detect any significant association between academic performance and residents' perceptions about their subjective quality of life. These findings might suggest that any knowledge gained, occurred independently and was unrelated to how people perceived their quality of life.

However, in contrast to the study done by De Oliveira Filho and Vieira (2007), this study explored the neurocognitive status of children, and it was found that two subscales had a significant relationship with academic achievement. These two subscales were from the perspective of cognition and negative moods. Both of these subscales are closely related to perceptions of negativity and can thus be linked to perceptions of the ability in catching up with school-work and the frequency negative feelings in children, both of which have been shown in this study to be the main factors affecting the QoL and at the same time is able to influence the academic achievement.

Regarding the relationship between neurocognitive status and academic achievement as measured by the CANTAB<sup>®</sup> battery of tests, no significant relationship was found between executive function and the academic achievement of the children as measured by their examination results. Similarly, there was no statistically significant relationship between visual memory and the academic achievement in these children.

Executive function is one of the important factors in learning, and with IED it is possible to detect the ability of the children in

simplediscrimination, reversal learning and also experiential learning. Those abilities are very useful to improve the learning abilities of children. Similarly, PAL is useful in the assessment of visual memory, and any learning difficulties with visual memory in young children will often lead to a significant problem in both long-term and short-term memory and difficulty in new learning. It also affects a person's ability to relearn appropriate behaviour (Cusimano, 2001). However, there is no association between both of the executive function and visual memory with the academic achievement. The main purpose of the current educational system is to measure he ability and success of students. These measurements still do not reflect the neurocognitive function of those children and some problems with neurocognition remain undetected. In Malaysia itself, the numbers of children with learning disabilities have almost doubled over the last three years period for primary school (473 to 656). This is an estimated increase of about 57% in the number of programs for special education for primary school level (Taib, 2008). In order to make academic achievement more efficient, the education system needs a neurocognitive profiling system that includes all the information about the quality of life neurocognitive wellness from the and neurocognitive status of the children.

# 5 CONCLUSION CATIONS

In conclusion, The fact that cognitive status of the children has no correlation with the academic achievement indicates that the school examination for the Standard one children cannot represent the cognitive ability regarding executive function and the visual memory. It can be explained that the academic achievement from the school still does not reflect the neurocognitive status of those children and some problems with neurocognition remain undetected. This information about the cognitive status of children is very important to improve the education system. Thus, the neurocognitive status can act as important indicators of a childs' academic achievement. This study will be expanded through the perspective of the children about other domain in cognitive status and its relation with their academic achievement.

### ACKNOWLEDGEMENTS

This study was supported by BRAINetwork Centre for Neurocognitive Science School of Health Universiti Sains Malaysia. The authors would like to express their deepest gratitude to children and teachers who participated in this study and the staff at primary schools in SK Zainab 2 Kota Bharu, Kelantan.

### REFERENCES

- Arnould, C., Penta, M., Renders, A., and Thonnard, J. L. (2004). ABILHAND-Kids A measure of manual ability in children with cerebral palsy. *Neurology*, 63(6): 1045-1052.
- Barnett, J. H., Blackwell, A. D., Sahakian, B. J., & Robbins, T. W. (2015). The paired associates learning (PAL) test: 30 years of CANTAB translational neuroscience from laboratory to bedside in dementia research. In *Translational Neuropsychopharmacology*, 449-474. Springer, Cham.
- Bernard, B. A., Stebbins, G. T., Siegel, S., Schultz, T. M., Hays, C., Morrissey, M. J., ... and Goetz, C. G. (2009). Determinants of quality of life in children with Gilles de la Tourette syndrome. *Movement disorders*, 24(7): 1070-1073.
- Breslow, L. (1999). New research points to the importance of using active learning in the classroom. *TLL Library*, 13 (1).
- Bullinger, M., Brütt, A. L., Erhart, M., Ravens-Sieberer, U., and BELLA Study Group. (2008). Psychometric properties of the KINDL-R questionnaire: results of the BELLA study. *European child & adolescent psychiatry*, 17(1): 125-132.
- Cella, D. F., and Tulsky, D. S. (1990). Measuring quality of life today: methodological aspects. *Oncology* (*Williston Park, NY*), 4(5): 29-38.
- Cognition, C. (2007). Cambridge Neuropsychological Tests Automated Battery (CANTAB), Eclipse Version. Cambridge, United Kingdom: Cambridge Cognition.
- Collinson, S. L., Tong, S. J., Loh, S. S., Chionh, S. B., and Merchant, R. A. (2014). Midlife metabolic syndrome and neurocognitive function in a mixed Asian sample. *International psychogeriatrics*, 26(8): 1305-1316.
- Cusimano, A. (2001). *Learning Disabilities--there is a Cure: A Guide for Parents, Educators, and Physicians*. Learning Disabilities.
- De Luca, C. R., Wood, S. J., Anderson, V., Buchanan, J. A., Proffitt, T. M., Mahony, K., and Pantelis, C. (2003). Normative data from the CANTAB. I: development of executive function over the lifespan. *Journal of clinical and experimental neuropsychology*, 25(2): 242-254.
- De Oliveira Filho, G. R., and Vieira, J. E. (2007). The relationship of learning environment, quality of life, and study strategies measures to anesthesiology

resident academic performance. Anesthesia & Analgesia, 104(6): 1467-1472.

- De Rover, M., Pironti, V. A., McCabe, J. A., Acosta-Cabronero, J., Arana, F. S., Morein-Zamir, S., ... and Sahakian, B. J. (2011). Hippocampal dysfunction in patients with mild cognitive impairment: a functional neuroimaging study of a visuospatial paired associates learning task. *Neuropsychologia*, 49(7): 2060-2070.
- Felce, D., and Perry, J. (1995). Quality of life: Its definition and measurement. *Research in developmental disabilities*, 16(1): 51-74.
- Heinzel, A., Northoff, G., Boeker, H., Boesiger, P., and Grimm, S. (2010). Emotional processing and executive functions in major depressive disorder: dorsal prefrontal activity correlates with performance in the intra–extra dimensional set shift. Acta neuropsychiatrica, 22(6): 269-279.
- Hofstede, G. (1984). The cultural relativity of the quality of life concept. *Academy of Management review*, 9(3): 389-398.
- Kamphuis, R., Theunissen, N., & Witb, J. (1997). Short communication health-related quality of life measure for children-the TACQOL. *J Appl Therapeut*, 1: 357-60.
- Kolsteren, M. M., Koopman, H. M., Schalekamp, G., & Mearin, M. L. (2001). Health-related quality of life in children with celiac disease. *The Journal of pediatrics*, 138(4): 593-595.
- Luciana, M. (2003). Practitioner review: computerized assessment of neuropsychological function in children: clinical and research applications of the Cambridge Neuropsychological Testing Automated Battery (CANTAB). Journal of Child Psychology and Psychiatry, 44(5): 649-663.
- Lukowski, A. F., Koss, M., Burden, M. J., Jonides, J., Nelson, C. A., Kaciroti, N., ... and Lozoff, B. (2010). Iron deficiency in infancy and neurocognitive functioning at 19 years: evidence of long-term deficits in executive function and recognition memory. *Nutritional neuroscience*, 13(2): 54-70.
- Lythe, K. E., Anderson, I. M., Deakin, J. F. W., Elliott, R., and Strickland, P. L. (2005). Lack of behavioural effects after acute tyrosine depletion in healthy volunteers. *Journal of Psychopharmacology*, 19(1): 5-11.
- O'Connell, H., Coen, R., Kidd, N., Warsi, M., Chin, A. V., and Lawlor, B. A. (2004). Early detection of Alzheimer's disease (AD) using the CANTAB paired Associates Learning Test. *International Journal of Geriatric Psychiatry*, 19(12): 1207-1208.
- Robbins, T. W., James, M., Owen, A. M., Sahakian, B. J., McInnes, L., and Rabbitt, P. (1994). Cambridge Neuropsychological Test Automated Battery (CANTAB): a factor analytic study of a large sample of normal elderly volunteers. *Dementia and Geriatric Cognitive Disorders*, 5(5): 266-281.
- Sahakian, B. J., and Owen, A. M. (1992). Computerized assessment in neuropsychiatry using CANTAB: discussion paper. *Journal of the Royal Society of Medicine*, 85(7): 399.

- Sharma, A. (2013). Cambridge Neuropsychological Test Automated Battery. In *Encyclopedia of Autism Spectrum Disorders*, pp. 498-515. Springer, New York, NY.
- Taib, M. N. M. (2008). School management concerning collaboration with social resources in community: Its' approaches and problems. *Primary and Middle Years Educator*, 6(2): 9-12.
- Verrips, E. G. H., Vogels, T. G. C., Koopman, H. M., Theunissen, N. C. M., Kamphuis, R. O. B. P., Fekkes, M., Wit, J. A. N. M. and Vanhorick, S. P. V. (1999). Measuring health-related quality of life in a child population. *The European Journal of Public Health*, 9(3):188-193.
- Vogels, A. G. C., Bruill, J., Stuifbergen, M., Koopman, H. M., and Verrips, G. H. W. (1999). Validity and reliability of a generic health-related quality of life instrument for adolescents, the TACQOL. *Quality of Life Research*, 630-630.
- Vogels, T., Verrips, G. H. W., Verloove-Vanhorick, S. P., Fekkes, M., Kamphuis, R. P., Koopman, H. M., ... and Wit, J. M. (1998). Measuring health-related quality of life in children: the development of the TACQOL parent form. *Quality of life research*, 7(5): 457-465.
- Vogels, T., Verrips, G. H. W., Koopman, H. M., Theunissen, N. C. M., Fekkes, M., & Kamphuis, R. P. (2000). TACQOL manual: parent form and child form. *Leiden: Leiden Center for Child Health and Pediatrics LUMC-TNO*.