

# Cell Cycle Arrest Activity of Alkaloid Fraction of *Litsea cubeba* Lour. Heartwoods towards HeLa Cancer Cell

Aminah Dalimunthe<sup>1\*</sup>, Poppy Anjelisa Zaitun Hasibuan<sup>1</sup>, Denny Satria<sup>2</sup>

<sup>1</sup>Department of Pharmacology, <sup>2</sup>Department of Pharmaceutical Biology  
Faculty of Pharmacy, University of Sumatera Utara, Medan, 20155

\*Presenting author

Keywords: Cell cycle, *Litsea cubeba* Lour. heartwood, alkaloid fractions, HeLa.

Abstract: Cervical cancer therapy with chemotherapeutic agents is limited because of drug resistance problem and toxic effect on normal tissue leads to immunosuppression and cardiotoxicity. This study was to investigated cell cycle arrest activity towards HeLa cell lines of *Litsea cubeba* Lour. heartwood alkaloid fraction. *Litsea cubeba* Lour. heartwood powder was extracted by maceration method with ethanol 96% and fractionated with n-hexane and chloroform at pH 3,7 and 9. The cytotoxic study was using MTT method and analysis cell cycle was using flow cytometry method. The IC<sub>50</sub> of ethanol extract, n-hexane and chloroform fractions at pH 3,7 and 9 at were 156.24 ± 2.96; 67.23 ± 0.63; 175.92 ± 2.40; 52.46 ± 0.34; and 94.81 ± 2.16 µg/mL respectively. The chloroform fractions at pH 7 concentration 25 and 10 µg/mL were caused accumulation in G<sub>2</sub>-M phase (33.84 and 29.08%). The results reveal that *Litsea cubeba* Lour. heartwood alkaloid fraction provides effective as cell cycle arrest. Our further study is to assess the mechanism of alkaloid fraction in inhibit metastasis in cervical cancer.

## 1 INTRODUCTION

Cancer is one of the high incidence dangerous diseases in human and presently there is a considerable number of new anticancer agents from natural products (Sharma, et al., 2011). According to WHO data, cancer is one of the leading cause of death worldwide especially cervical cancer (Berrington and Lall, 2012). Cervical cancer therapy with chemotherapeutic agents is limited cause of drug resistance and toxic side effect on normal tissue leads to some effects such as immunosuppression and cardiotoxicity (Jemal, et al., 2010; Tyagi, et al., 2004).

Attarasa (*Litsea cubeba* (Lour.) is a plant from Lauraceae family which contain many essential oils which used as antidepressants, antiinflammation,

antioxidant, pesticide, antimicrobial, anticancer on breast cancer and neuro pharmacology. The methanol extract from attarasa fruits showed to be active on HeLa cell lines which cause apoptosis through activation of caspase 3/7 (Trisonthi, et al., 2014; Piyapat, et al., 2013). There are more than forty isoquinoline alkaloids that contained in *Litsea* genus which are active as antibacterial agents against *Staphylococcus aureus* (Feng, et al., 2009). The heartwoods of *Litsea cubeba* contained high level of phenolic and flavonoid and found to be active as antioxidant and has anti breast cancer activity which causes cell cycle inhibition. Alkaloids compound which isolated from heartwood have antioxidant activity with DPPH and ABTS methods (Dalimunthe, et al., 2016; Dalimunthe, et al., 2017; Dalimunthe, et al., 2018). The aim of this study was

to assess cell cycle arrest activity of alkaloid fraction of *Litsea cubeba* Lour. heartwoods on HeLa cells.

## 2 MATERIALS AND METHODS

### 2.1 Fractions Preparation

Fresh heartwoods of *Litsea cubeba* Lour. was collected from Balige subdistrict, Sumatera Utara province, Indonesia. The air-dried and powdered heartwoods of *Litsea cubeba* (Lour.) (1 kg) were repeatedly macerated with ethanol 96% (3x3 d, 7.5 L), The filtrate was evaporated to give a viscous extract. Viscous extract was fractionated with n-hexane and continue with chloroform at pH 3,7 and 9 (Rosidah, et al., 2018; Dalimunthe, et al., 2018; Satria, et al., 2015).

### 2.2 Cytotoxicity Assay

Extract and alkaloid fractions were submitted for cytotoxicity test. In that way, HeLa cell line was grown in RPMI medium containing 10% Fetal Bovine Serum (Gibco), 1% penicillin-streptomycin (Gibco), and fungizone 0.5% (Gibco) in a flask in a humidified atmosphere (5% CO<sub>2</sub>) at 37°C. The inoculum seeded at 1 x 10<sup>4</sup> cells/mL at an optimal volume of 0.1 mL per well. After 24 h incubation, the medium was discharged and treated by fractions. After incubation for 24 h, the cells were incubated with 0.5 mg/mL MTT for 4 h at 37°C. Viable cells reacted with MTT to produce purple formazan crystals. After 4 h, SDS 10% as stopper (Sigma) in 0.01N HCl (Merck) was added to dissolve the formazan crystals. The cells were incubated for 24 h in room temperature and protected from light. After incubation, the cells were shaken, and absorbance was measured using microplate reader at  $\lambda$  595 nm.

The data which were absorbed from each well were converted to percentage of viable cells (Hasibuan, et al., 2015 and Nurrochmad, et al., 2014).

### 2.3 Cell Cycle Inhibition Assay

HeLa cells (7.5x10<sup>5</sup> cells/well) were seeded into 6-well plate and incubated for 24 h. After that, the cells were treated and then incubated for 24 h. Both floating and adherent cells were collected in conical tube using trypsin 0.025%. The cells were washed thrice with cold PBS and centrifuged at 2500 rpm for 5 min. The supernatant was separated, while the sediment was collected and fixed in cold 70% ethanol in PBS at 4°C for 1 h. The cells were washed thrice with cold PBS and resuspended then centrifuged at 3000 rpm for 3 min and PI kit (containing PI 40  $\mu$ g/mL and RNase 100  $\mu$ g/mL) added to sediment and resuspended and incubated at 37°C for 30 min. The samples were analyzed using FACScan flow cytometer. Based on DNA content, percentage of cells in each of stage in cell cycle (G1, S and G2/M) were calculated using ModFit Lt. 3.0.s (Harahap, et al., 2018 and Satria, et al., 2017).

### 2.4 Statistical Analysis

The results were presented as means  $\pm$  SD.

## 3 RESULTS

### 3.1 Inhibitory Concentration 50% (IC<sub>50</sub>)

MTT method was used to determine cell viability after incubation for 24 h. In every treatment extract and alkaloid fractions were shown in Table 1.

Table 1. IC<sub>50</sub> value of extract and alkaloid fractions of *Litsea cubeba* heartwood with MTT assay (Mean  $\pm$  SD, 3 times of replication)

Treatment	IC <sub>50</sub> ( $\mu$ g/mL)
Ethanol Extract	156.24 $\pm$ 2.96
n-hexane Fraction	67.23 $\pm$ 0.63
Chloroform Fraction pH 3	175.92 $\pm$ 2.40
Chloroform Fraction pH 7	52.46 $\pm$ 0.34
Chloroform Fraction pH 9	94.81 $\pm$ 2.16
Cisplatin	24.01 $\pm$ 0.31

### 3.2 Effect on Cell Cycle

To evaluate the effect of chloroform fraction at pH 7 (CF-7) to increase cell death by modulating cell cycle, we concentrated on it for further studies using flow cytometry method.

The effect of CF-7 at 25 and 10  $\mu\text{g/mL}$  is given in Figure 1. Whereas treatment of CF-7 at 25 and 10  $\mu\text{g/mL}$  caused cell accumulation at  $G_2/M$  phase (33.84% and 29.08%) and for control cell (17.78%).




Figure 1. Percentage of cell cycle phase of HeLa cells were treated for 24h. (a) Control cell, (b) 25  $\mu\text{g/mL}$  ( $1/2\text{ IC}_{50}$ ), (c) 10  $\mu\text{g/mL}$  ( $1/5\text{ IC}_{50}$ ).

## 4 DISCUSSION

The cytotoxicity estimate of herbal is correlated to content of active compound in these plants including *Litsea cubeba* Lour. Alkaloids as major compound have main role in cytotoxicity effect (Yadav, et al., 2010). *Litsea* genus is rich in isoquinoline alkaloids and for *Litsea cubeba* Lour has been found two alkaloid (+)-*N*-(methoxy-carbonyl) *N*-norlaurosoltzine and (+)-*N*-(methoxy-carbonyl)-*N*-norglaucine (Feng, et al., 2009). Alkaloids are the compound which potentially in inhibits the cancer proliferation for the example berberine is an isoquinoline alkaloid which inhibits proliferation of multiple cancer cell line by inducing cell cycle arrest at  $G_0/G_1$  or  $G_2/M$  phases and by apoptosis (Sun, et al., 2009; Eom, et al., 2010; Burgeiro, et al., 2011). Inhibition of tumor invasion and metastasis is the mechanism of action of berberine (Tang, et al., 2009; Ho, et al., 2009). Evodiamine is a quinolone alkaloid inhibits topoisomerase enzyme, induces DNA damage, exhibit  $G_2/M$  phase arrest (Liao, et al., 2005; Kan, et al., 2004; Huang, et al., 2004).

## ACKNOWLEDGEMENTS

We gratefully thank to Research Center University of Sumatera Utara through Hibah Talenta “Hibah Penelitian Dasar” Research Grant 2018 “No:2590/UN5.1.R/PPM/2018” for financial support in the study.

## REFERENCES

- Berrington D and Lall N (2012). Anticancer activity of certain herbs and spices on the cervical epithelial carcinoma (HeLa) cell line. eCAM, Article ID 564927.
- Burgeiro Z, Gajate C, Dakir EH, Villa-Pulgarín JA, Oliveira PJ. (2011). Involvement of mitochondrial and B-RAF/ERK signaling pathways in berberine-induced apoptosis in human melanoma cells. *Anti-Cancer Drugs*, 22(6), 507–18.
- Dalimunthe A, Achmad S, Satria D. (2016). Phenolic, flavonoid content and antioxidant activities of ethylacetate extract of *litsea cubeba* (lour.) pers. barks. *Der Pharma Chemica*, 8, 466-468.
- Dalimunthe A, Hasibuan PAZ, Satria D. (2017). Cell cycle arrest activity of *Litsea cubeba* Lour. heartwood

- and fruit extracts against T47D breast cancer cells. *Asian J Pharm Clin Resc*, 10(11), 404-6.
- Dalimunthe A, Hasibuan PAZ, Silalahi J, Sinaga SF, Satria D. (2018). Antioxidant activity of alkaloid compounds from *Litsea cubeba* Lour. *Orient J Chem*, 34(2), 1149-52.
- Eom KS, Kim HJ, So HS, Park R, Kim TY. (2010). Berberine-induced apoptosis in human glioblastoma T98G Cells Is mediated by endoplasmic reticulum stress accompanying reactive oxygen species and mitochondrial dysfunction. *Biol Pharm Bull*, 33(10), 1644-9.
- Feng T, Rong-Ting Z, Qin-Gang T, Xiang-Yun Z. (2009). Two new isoquinoline alkaloids from *Litsea cubeba*. *Z Naturforsch*, 64, 871-4.
- Harahap U, Hasibuan PAZ, Sitorus P, Arfian N, Satria D. (2018). Antimigration activity of an ethylacetate fraction of *Zanthoxylum acanthopodium* DC. fruits in 4T1 breast cancer cells. *Asian Pac J Cancer Prev*, 19(2), 565-9.
- Hasibuan PAZ, Harahap U, Sitorus P, Satria D (2016). Ethylacetate extract of *Zanthoxylum acanthopodium* DC. fruit against doxorubicin-resistance T47D cells. *Der Pharma Chemica*, 8(20), 172-174.
- Hasibuan PAZ, Jessy C, Denny S (2015). Combination effect of ethylacetate extracts of *Plectranthus ambonicus* (Lour.) Spreng. with doxorubicin against T47D breast cancer cells. *Int J Pharm Pharm Sci*, 7, 155-9.
- Ho YT, Yang JS, Li TC. (2009). Berberine suppresses *in vitro* migration and invasion of human SCC-4 tongue squamous cancer cells through the inhibitions of FAK, IKK, NF- $\kappa$ B, u-PA and MMP-2 and -9. *Cancer Letters*, 279(2), 155-62.
- Huang YC, Guh JH, Teng CM. (2004). Induction of mitotic arrest and apoptosis by evodiamine in human leukemic T lymphocytes. *Life Sciences*, 75(1), 35-49.
- Jemal A, Siegel R, Xu J, Ward E (2010). *CA Cancer J Clin*.
- Kan SF, Huang WJ, Lin LC, Wang PS. (2004). Inhibitory effects of evodiamine on the growth of human prostate cancer cell line LNCaP. *Int J Canc*, 110(5), 641-51.
- Liao CH, Pan SL, Guh JH. (2005). Antitumor mechanism of evodiamine, a constituent from Chinese herb *Evodiae fructus*, in human multiple-drug resistant breast cancer NCI/ADR-RES cells *in vitro* and *in vivo*. *Carcinogenesis*, 26(5), 968-75.
- Nurrochmad A, Lukitaningsih E, Meiyanto E (2011). Anti cancer activity of rodent tuber (*Thyphonium flagelliforme* (Lodd.) Blume) on human breast cancer t47d cells. *International Journal of Phytomedicine*, 3, 138-146.
- Piyapat T, Miyagawa K, Tamura H. (2013). Induction of apoptosis in HeLa cells by methanol extract of *Litsea cubeba* fruit residue from essential oil extraction. *J Life Sci*. 7(9), 928-34.
- Rosidah, Hasibuan PAZ, Haro G, Puteri M, Satria D. (2018). Antioxidant activity of alkaloid fractions of *Zanthoxylum acanthopodium* DC. fruits with 1,1-diphenyl-2-picrylhydrazyl assay. *Asian J Pharm Clin Res*, 11(1), 33-4.
- Satria D, Furqan M, Hadisahputra S, Rosidah (2015). Combinational effects of ethylacetate extract of *Picria fel-terrae* Lour. and doxorubicin on T47D breast cancer cells. *Int J Pharm Pharm Sci*, 7, 73-6.
- Satria D, Silalahi J, Haro G, Ilyas S, Hasibuan PAZ. (2017). Antioxidant and antiproliferative activities of an ethylacetate fraction of *Picria fel-terrae* Lour. herbs. *Asian Pac J Cancer Prev*, 18(2), 399-403.
- Sharma J, Pitchaiah G, Satyawati D, Rao JV, Kumar-Vikram HS (2011). *In vitro* anticancer activity of methanolic extract of roots of *Glochidion zeylanicum* Gaertn. *Int J Res Pharmaceut Biomed Sci*, 2, 760-764.
- Sun Y, Xun K, Wang Y, Chen X. (2009). A systematic review of the anticancer properties of berberine, a natural product from Chinese herbs. *Anti-Cancer Drugs*, 20(9), 757-69.
- Tang F, Wang D, Duan C. (2009). Berberine inhibits metastasis of nasopharyngeal carcinoma 5-8F cells by targeting rho kinase-mediated ezrin phosphorylation at threonine 567. *J Bio Chem*, 284(40), 27456-66.
- Trisonthi P, Akihiko S, Hisashi N, Hirotooshi T. (2014). A new diterpene from *Litsea cubeba* fruits: structure elucidation and capability to induce apoptosis in HeLa cells. *Molecules*, 19, 6838-50.
- Tyagi AK, Agarwal C, Chan DCF, Agarwal R (2004). *Oncology Reports*, 11, 493-499.
- Yadav VR, Sahdeo P, Bokyung S, Ramaswamy K, Bharat BA (2010). Targeting inflammatory pathways by triterpenoids for prevention and treatment of cancer. *Toxins*, 2, 2428-66.
- Zihlif M, Afifi F, Abu-Dahab R, Majid A, Somrain H, Saleh M (2013). The antiangiogenic activities of ethanolic crude extracts of four *Salvia* species. *Complementary & Alternative Medicine*, 13, 358-65.