

A U-Net-Based Temperature Bias Correction Method for the REMO2015 Regional Climate Model in CORDEX-EA

Shibin Zheng¹, Chenwei Shen² and Bin Li²

¹*School of Computer and Artificial Intelligence, Zhengzhou University, Zhengzhou, China*

²*Dawning Information Industry Company Limited, Beijing, China*

Keywords: Bias Correction, CORDEX East Asia, Deep Learning, U-Net.

Abstract: Regional climate models suffer from insufficient resolution and deficiencies in their dynamic processes, leading to systematic biases in surface air temperature simulations that require correction. In this research, a deep learning bias correction model, CE-MS-Unet, is proposed. This model incorporates multi-scale residual blocks and calendar month data to improve surface air temperature simulations of the REMO2015 regional climate model during the second phase of the Coordinated Regional Downscaling Experiment East Asia (CORDEX-EA-II) over mainland China. Experimental results indicate that, compared to Linear Scaling, Quantile Delta Mapping, and the deep learning model CU-net, CE-MS-Unet performs better in correcting climate averages and seasonal cycles, resulting in corrected data with greater overall agreement and improved spatial correlation. It effectively reduces biases and provides more accurate climate predictions. This study offers new insights and methods to improve the bias correction of temperature in regional climate models.

1 INTRODUCTION

In the field of climatology, Global Climate Models (GCMs), which couple global atmospheric, oceanic, and terrestrial systems, are widely used for studying long-term climate change and future climate projections. However, the relatively low grid resolution of GCMs limits their capacity to accurately capture climate changes on a regional scale. The application of dynamically downscaled Regional Climate Models (RCMs) driven by GCMs within a region can provide higher-resolution local information, thereby enhancing the accuracy of detailed climate impact assessments (Giorgi et al., 1999). The Coordinated Regional Climate Downscaling Experiment (CORDEX), launched by the World Climate Research Programme (WCRP), provides high-resolution regional climate projections for land areas inhabited by most of the global population using multiple RCMs (Gutowski et al., 2016). This study focuses on CORDEX-East Asia (CORDEX-EA), the East Asian branch of the CORDEX program. Previous studies indicate that the RCMs used in the CORDEX-EA-II experiments can effectively simulate and project surface air temperature and precipitation (Yu et al., 2020).

However, due to the inherent limitations in dynamical processes and physical parameterization within RCMs, as well as biases inherited from their driving GCMs, the simulated outputs still have considerable systematic biases. Statistical bias correction methods are commonly used to reduce biases and improve the accuracy of future climate projections. These methods establish a statistical relationship between simulated and observed data to minimize their distributional differences. Two widely used techniques are Linear Scaling (LS) and Quantile Delta Mapping (QDM). LS adjusts the mean or standard deviation of data through a simple linear transformation and efficiently corrects seasonal temperature variations (Chen et al., 2022). However, it assumes the correction factor remains valid under future climate conditions, which can lead to inaccuracies as the climate changes. QDM, an advanced version of Quantile Mapping (QM), corrects both the distribution and trends of simulated data by mapping quantile changes while retaining the model's predicted climate change signals. (Tong et al., 2021). Nevertheless, QDM is less effective at managing spatial correlations and intermittency.

In recent years, deep learning models have been increasingly utilized in meteorology, resulting in the development of numerous artificial neural network-based bias correction methods (de Burgh-Day et al.,

2023). Originating from computer vision, these models treat meteorological bias correction as a regression task for fitting image features, using raw data as input predictors for training. Several bias correction methods based on Generative Adversarial Networks (GANs) have been proposed. GANs can be trained on unpaired image data to learn the bias distribution of GCMs and generate corrected images, making them naturally effective for adjusting GCM outputs without corresponding observational data and capturing spatial precipitation patterns (Pan et al., 2021; Hess et al., 2023). Additionally, convolutional neural network (CNN)-based methods that are widely used in short-term weather forecasting have shown their potential in climate model bias correction and downscaling (Sha et al., 2020). CNN-based models extract multi-scale spatial features through convolutional and pooling layers, use multi-channel input data to capture complex nonlinear relationships between different variables, thereby potentially improving the bias correction performance of GCMs or RCMs (Kesavavarthini et al., 2023; Wang and Tian, 2022). Recently, the U-net, a CNN derivative originally developed for medical image segmentation, has also been applied to meteorological bias correction (Molina et al., 2023). With its encoder-decoder structure, U-net can effectively extract features and restores spatial information. Compared to traditional CNNs, it captures multi-scale spatial details while producing outputs that match the original image size.

Although previous work on bias correction for RCMs in the CORDEX-EA experiments has primarily employed traditional statistical methods, no studies have explored deep learning-based correction approaches (Chen et al., 2022; Tong et al., 2021). To improve surface air temperature simulations of regional climate models in the CORDEX-EA-II experiments over mainland China, this study implements a deep learning bias correction model based on U-net. The choice to forgo a GAN-based approach was driven by two main reasons: first, the large data requirements of GANs are challenging to meet given that the CORDEX-EA experiment's simulations span only up to 35 years; and second, the instability and convergence challenges inherent in the GAN's architecture complicates its application and training (Yu et al., 2024). This research introduces a new CE-MS-Unet model that incorporates multi-scale residual blocks and one-hot encoding of calendar month data. When applied to surface air temperature bias correction in the REMO2015 regional climate model, this model achieves better overall agreement and more accurate temperature

cycle correction compared to traditional methods and the CU-net model. Consequently, it can support more reliable long-term regional surface air temperature predictions.

The paper is organized as follows: Section 2 details the study area and data preprocessing steps. Section 3 describes the implemented bias correction methods, including two statistical and two deep learning approaches. Section 4 covers the experimental setup and analyzes the results, while Section 5 concludes with a summary.

2 STUDY AREA AND DATA

As shown in Figure 1, this study focuses on a region from the CORDEX-EA-II experiment that primarily covers mainland China, extending from 18°N to 55°N and from 75°E to 135°E. To further evaluate the performance of various bias correction methods at a smaller spatial scale, five subregions within the study area were selected.




Figure 1: Topography of the study area and its five subregions: Southern China (SC), Northern China (NC), Northeastern China (NE), Northwestern China (NW), and the Tibetan Plateau (TP).

The bias correction uses RCM output data from REMO2015, developed by the Climate Service Center Germany (GERICS). TAS, TASMAL and TASMAL were selected from the historical simulation data of CORDEX-EA-II experiment. Additionally, digital elevation model (DEM) data were included. These features are related to air temperature within the climate system, which can improve the accuracy of the deep learning model in correcting temperature biases (Zhang et al., 2022). The Asian Precipitation-Highly-Resolved Observational Data Integration

Towards the Evaluation of Water Resources (APHRODITE, abbreviated as APHRO) gridded dataset(V1101) was used as reference data. Detailed information about datasets is provided in Table 1.

Table 1: Datasets used in this study.

Datasets	Variables used	Temporal
APHRO DITE	Daily Mean Temperature (TAVE)	1971- 2005
SRTM	Digital Elevation Model (DEM)	1971- 2005
REMO2 015 Output	Near-Surface Air Temperature (TAS)	1971- 2005
	Daily Minimum Near- Surface Air Temperature (TASMIN)	
	Daily Maximum Near- Surface Air Temperature (TASMAX)	

Bilinear interpolation was applied to resample the meteorological variables from REMO2015 to align with the $0.25^\circ \times 0.25^\circ$ resolution of the APHRO dataset. For deep learning methods, data from 1971 to 2000 were used for training and validation, while data from 2001 to 2005 served as the test set. To enable the model to learn temperature variation patterns across different climate states, a strategy similar to Pan et al. (2021) was employed: from 1971, the first four years of each five-year period were included in the training set, with the final year in the validation set. For each time step T within these datasets, reference data from the same month within a five-year window around T were randomly selected as the target data. All meteorological variables were standardized using Z-score normalization.

3 METHODS

This study implemented two widely used statistical methods and two U-net based deep learning methods. LS and QDM were selected as baseline statistical methods for the CE-MS-Unet model, while the CU-net model was used as the baseline for the deep learning methods.

3.1 Linear Scaling

Linear Scaling aims to minimize the mean bias between RCM predictions and observational data over monthly time series (Teutschbein and Seibert,

2012). An additive scaling approach is used to compute the corrected value of meteorological variable X at time step i :

$$X_{bc,p}(i) = X_{sim,p}(i) + \mu_m(X_{obs,c}(i)) - \mu_m(X_{sim,c}(i)) \quad (1)$$

Where $\mu_m(X_{...}(i))$ is the long-term monthly average temperature for the month corresponding to time step i . In the subscripts, sim denotes the RCM simulated value, obs the observed value, bc the bias-corrected value, p the scenario period, and c the control period.

3.2 Quantile Delta Mapping

Quantile Delta Mapping is a technique used to correct distributional biases between RCM predictions and observational data. Unlike the conventional Quantile Mapping method, QDM not only matches RCM data with observational data during the control period but also accounts for changes between the control period and the scenario period (Tong et al., 2021).

Specifically, for the simulated climate variable X , the non-exceedance probability $\varepsilon(i)$ at time step i during the scenario period is first calculated:

$$\varepsilon(i) = F_{sim,p}(X_{sim,p}(i)) \quad (2)$$

Next, the bias-corrected value $X'_{bc,p}(i)$ is determined by substituting the non-exceedance probability into the inverse cumulative distribution function of the observational data from the control period:

$$X'_{bc,p}(i) = F_{obs,c}^{-1}[\varepsilon(i)] \quad (3)$$

The absolute change in quantiles between control period and scenario period is then calculated as:

$$\Delta(i) = F_{sim,p}^{-1}[\varepsilon(i)] - F_{sim,c}^{-1}[\varepsilon(i)] = X_{sim,p}(i) - F_{sim,c}^{-1}\{F_{sim,p}[X_{sim,p}(i)]\} \quad (4)$$

At the time step i during scenario period, the final corrected temperature is obtained by adding the absolute change amount to the bias-corrected value.

$$X_{bc,p}(i) = X'_{bc,p}(i) + \Delta(i) \quad (5)$$

3.3 CU-Net

Based on the study by Han et al. (2021), we introduce the CU-net model to correct the surface air temperature simulation biases of the REMO2015 regional climate model. CU-net is a deep learning model




Figure 2: Architecture of the CE-MS-Unet model. The model has two input layers: one for meteorological factors and the other for calendar month data.

designed for bias correction in meteorology, with an architecture similar to U-net. When meteorological data is fed into CU-net, the left half of its U-shaped structure, consisting of a CNN-based convolutional encoding module, automatically extracts high-level features from the data. The right half, which consists of an upsampling module, performs decoding operations to progressively restore the compressed feature maps to their original size. During this upsampling process, CU-net employs the "copy and concatenate" operation that merges feature maps from the encoder and decoder along the channel dimension.

CU-net differs from the original U-net in its use of sub-pixel convolution in the decoder. When applied to the expansion of meteorological feature maps, sub-pixel convolution enhances computational efficiency and reduces the loss of valuable information during image reconstruction.

3.4 CE-MS-Unet

Building upon the CU-net architecture, this study introduces multi-scale residual blocks and one-hot encoding of calendar month data, leading to a new model: the Calendar-Embedded Multi-Scale Residual U-net (CE-MS-Unet). Figure 2 illustrates the structure of CE-MS-Unet. CE-MS-Unet replaces the sequential convolutions in each layer with multi-scale residual blocks and incorporates calendar month data as additional input at the deepest layer.




Figure 3: Structure of the multi-scale residual block.

Biases in RCM temperature simulations may result from interactions between climate processes occurring at different spatial scales, such as local effects and large-scale weather systems. Therefore, more effectively capturing meteorological features across multiple spatial scales can potentially improve bias correction performance (Faijaroenmongkol et al., 2023). As shown in Figure 3, the Multi-Scale Residual Block captures multi-scale information in the temperature field using parallel convolutional kernels of different sizes. These multi-scale features are then fused and passed to the next network layer through a Residual Connection. The use of feature fusion and residual connections stabilizes deep network training, helping prevent overfitting and reduce noise and uncertainty in temperature data.

Additionally, the Exponential Linear Unit (ELU) activation function is used in all convolutional layers.

Temperature shows significant seasonal variations, with distinct patterns and characteristics across different months. The use of calendar data in deep learning models has been successfully applied to precipitation bias correction (Ling et al., 2022). To improve the model's ability to capture temperature bias characteristics across different months and seasons, calendar month data was used as an additional input. These data are represented as a 12-dimensional one-hot encoded vector, where each dimension corresponds to a month and is then fused with the feature maps at the model's deepest layer. Before fusion, learnable scaling factors are applied to dynamically adjust the weights of the two data sources, optimizing their relative influence during the fusion process. Introducing calendar month data at the deepest layer is intended to preserve the CU-net model's original spatial feature extraction capabilities while integrating temporal information with high-level abstract features, thereby enhancing the model's final correction output more effectively.

4 EXPERIMENT AND RESULTS

4.1 Training Setting

During training, the ADAM optimizer was used with an initial learning rate of 0.001 and a batch size of 16. The total number of epochs was set to 50. Dynamic learning rate adjustment were employed: if the validation loss did not decrease for two consecutive epochs, the learning rate was halved. After training, the model weights with the lowest validation loss were saved. Both models utilized a custom loss function that considers the Mean Squared Error (MSE) at each grid point, as well as the MSE of the overall data mean and standard deviation, defined as follows:

$$L = \text{MSE}(y_i - y'_i) + 3 \times \text{MSE}(y_{\text{mean}} - y'_{\text{mean}}) + 3 \times \text{MSE}(y_{\text{std}} - y'_{\text{std}}) \quad (6)$$

Where y_i and y'_i represent the observed and corrected values, with the subscripts mean and std denoting their mean and standard deviation, respectively.

Both deep learning models were implemented using TensorFlow 2.9 and Python 3.9 and were trained on four GPU-like accelerators. The accelerator adopts a GPU-like architecture consisting

of a 16GB HBM2 device memory and many compute units, with peak FP64 performance of 7.0TFLOPS.

4.2 Statistical Performance Metrics

To evaluate the effectiveness of each bias correction method, mean absolute error (MAE), root mean squared error (RMSE), and spatial correlation coefficient (SCC) were employed. MAE and RMSE is calculated as:

$$\text{MAE} = \frac{1}{n} \sum_{i=1}^n |y_i - y'_i| \quad (7)$$

$$\text{RMSE} = \sqrt{\frac{1}{n} \sum_{i=1}^n (y_i - y'_i)^2} \quad (8)$$

Where y_i is the observed values and y'_i is the corrected or original values. The Spatial Correlation Coefficient (SCC) is used to evaluate the correlation between the spatial distributions of temperature values before and after correction:

$$\text{SCC} = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sqrt{\sum_{i=1}^n (x_i - \bar{x})^2} \sqrt{\sum_{i=1}^n (y_i - \bar{y})^2}} \quad (9)$$

Where x_i and y_i represent the values in the observed and corrected spatial distributions, and \bar{x} and \bar{y} are their respective means.

4.3 Results

4.3.1 Overall Agreement

The overall agreement between the corrected and observed surface air temperature was evaluated using MAE and RMSE values calculated for each grid point across the entire study area and its five subregions. Detailed results are presented in Table 2. Across the study area, four corrected results exhibited different levels of improvement over the original RCM data. LS showed a slight advantage compared to QDM, whereas CU-Net and CE-MS-Unet outperformed LS. CE-MS-Unet performed the best, reducing the MAE and RMSE values by 0.23 and 0.24 respectively, compared to CU-Net. The four methods varied in performance across subregions. Among statistical methods, QDM outperformed LS in MAE and RMSE in the NW and TP regions, while LS performed better in the others. For deep learning methods, CE-MS-Unet consistently surpassed CU-Net across all regions. In four of the five subregions (excluding TP), deep learning methods showed better consistency than statistical methods, with CE-MS-Unet yielding

Table 2: MAE and RMSE values for RCM output and four bias-corrected results across the entire study area and its five subregions, the best-performing values are highlighted in bold.

Regions	MAEs					RMSEs				
	RCM	LS	QDM	CU-Net	CE-MS-UNet	RCM	LS	QDM	CU-Net	CE-MS-UNet
SC	3.24	2.97	3.08	2.18	1.94	4.21	3.89	4.08	2.89	2.58
NC	3.65	3.28	3.38	2.67	2.46	4.70	4.25	4.41	3.42	3.12
NE	3.97	3.69	3.68	3.01	2.66	4.99	4.62	4.69	3.75	3.27
NW	2.94	2.28	2.18	2.07	1.90	3.71	2.93	2.83	2.59	2.37
TP	4.38	2.37	2.09	2.75	2.59	5.63	3.09	2.71	3.69	3.45
Overall	4.58	3.93	4.01	3.88	3.65	6.05	5.24	5.41	5.19	4.95




Figure 4: Spatial-distribution of mean temperature biases for the testing period (2001-2005) from (a) RCM and (b-e) four bias-correction methods (unit: °C). The spatial average RMSEs (the upper one) and annual average daily map correlations (the lower one) between the RCM/corrected outputs and observations are provided in lower right corner of the panels.

the best results. In the TP region, the complex terrain results in larger biases in RCM simulations. Traditional methods process data in a relatively simple way, making them better suited to this scenario. In contrast, deep learning models struggle to learn temperature bias characteristics due to the large amount of high-error data. Consequently, QDM performs best in the TP region. These results suggest that, in terms of overall agreement with surface air temperature data, the two U-Net-based deep learning methods provide superior corrections across most regions, with CE-MS-UNet yielding the most consistent results.

4.3.2 Spatial Distribution Bias

As shown in Figure 4, the five-year average temperature bias between the corrected results and observational data was calculated to assess each method's ability to correct spatial biases. The RMSE of the original data's annual average temperature reached 2.41, while all four correction methods significantly reduced this bias, lowering the RMSE to below 1. Owing to their superior spatial feature extraction capability, CU-net and CE-MS-UNet not only effectively reduced the bias but also better preserved the original spatial patterns of the RCM.




Figure 5: Annual cycles of temperature biases from REMO2015 and four bias-correction results over five subregions.

CE-MS-Unet reduced the bias to below 1°C in most regions and eliminated the cold bias in high-latitude areas seen with LS and QDM, resulting in a more balanced cold-warm bias distribution.

Additionally, the spatial correlation coefficients (SCC) between the five-year annual average temperatures of each dataset and the observational data were calculated. The results indicated that the original RCM data had a SCC of 0.98, while all four correction methods improved it to 1. To further assess each method's ability to enhance spatial correlation, the approach of Wang and Tian (2022) was employed. This method flattens the 2D spatial data into a 1D vector to calculate daily map correlations, which are then averaged over the 5 years. Figure 4 indicates that CE-MS-Unet achieved the highest annual average daily map correlation. Although CU-net also demonstrated a relatively high map correlation, its RMSE was notably higher. Taking both metrics into account, CE-MS-Unet has a clear advantage in correcting spatial biases of temperature.

4.3.3 Temporal Skill

Figure 5 illustrates the regional monthly mean temperature biases between the corrected results and observational data. In four subregions excluding TP, CE-MS-Unet, LS, and QDM significantly reduced the monthly mean temperature bias, bringing it below 1°C for most months and closely matching the observational climatology. CU-Net reduced the bias in average temperatures for spring, summer, and autumn, but showed a substantial warm bias in winter. CE-MS-Unet effectively addressed the winter bias

issue observed in CU-Net and demonstrated comparable capabilities to LS and QDM across four subregions. Moreover, the deviations in the lowest and highest monthly mean temperatures corrected by LS and QDM were around 3°C, while those corrected by CE-MS-Unet were closer to 2°C, indicating that CE-MS-Unet had less variability than the traditional methods. In the TP region, both deep learning methods were less effective than LS and QDM in reducing the significant cold bias in RCM simulations. This result aligns with the overall agreement section and is attributed to higher errors and lower data quality in the region's simulations.

5 CONCLUSIONS

To improve the accuracy of surface air temperature simulations from the REMO2015 model within the CORDEX-EA project over mainland China, we presented a U-Net-based bias correction model, CE-MS-Unet. Experimental results demonstrate that, compared to traditional statistical methods like Linear Scaling and Quantile Delta Mapping, as well as the existing deep learning model CU-net, CE-MS-Unet is better at capturing the spatial and temporal features of surface air temperature. This improvement is achieved by incorporating multi-scale residual blocks and embedding calendar month data. In East Asia, CE-MS-Unet excels in reducing MAE and RMSE, while also providing superior correction for spatial distribution and seasonal cycles. Although slightly inferior to QDM in the Tibetan Plateau, CE-MS-Unet

overall outperforms LS, QDM, and CU-net in correcting spatial and temporal biases in REMO2015's surface air temperature simulations.

Future work could explore further adjustments to the CE-MS-Unet structure, such as integrating attention mechanisms, designing more sophisticated methods for calendar data fusion, and enhancing the model's bias correction performance in the Tibetan Plateau. Ablation studies could also be conducted to improve the model's interpretability. Additionally, testing CE-MS-Unet's performance in CORDEX experiments outside East Asia would help validate its generalization and applicability.

ACKNOWLEDGEMENTS

This work was supported by the State Key RandD Program of China (No. 2021YFB0300200).

REFERENCES

- Giorgi, F., and Mearns, L. O. (1999). Introduction to special section: Regional climate modeling revisited. *Journal of Geophysical Research: Atmospheres*, 104(D6), 6335-6352.
- Gutowski Jr, W. J., Giorgi, F., Timbal, B., Frigon, A., Jacob, D., Kang, H.-S., Krishnan, R., Lee, B., Lennard, C., and Nikulin, G. (2016). WCRP coordinated regional downscaling experiment (CORDEX): a diagnostic MIP for CMIP6.
- Yu, K., Hui, P., Zhou, W., and Tang, J. (2020). Evaluation of multi-RCM high-resolution hindcast over the CORDEX East Asia Phase II region: Mean, annual cycle and interannual variations. *International Journal of Climatology*, 40(4), 2134-2152.
- Chen, J., Yang, Y., and Tang, J. (2022). Bias correction of surface air temperature and precipitation in CORDEX East Asia simulation: What should we do when applying bias correction?. *Atmospheric Research*, 280, 106439.
- Tong, Y., Gao, X., Han, Z., Xu, Y., Xu, Y., and Giorgi, F. (2021). Bias correction of temperature and precipitation over China for RCM simulations using the QM and QDM methods. *Climate Dynamics*, 57, 1425-1443.
- de Burgh-Day, C. O., and Leeuwenburg, T. (2023). Machine learning for numerical weather and climate modelling: a review. *Geoscientific Model Development*, 16(22), 6433-6477.
- Kesavavarthini, T., Rajesh, A. N., Venkata Srinivas, C., and Kumar, T. L. (2023). Bias correction of CMIP6 simulations of precipitation over Indian monsoon core region using deep learning algorithms. *International Journal of Climatology*, 43(8), 3749-3767.
- Molina, M. J., O'Brien, T. A., Anderson, G., Ashfaq, M., Bennett, K. E., Collins, W. D., Dagon, K., Restrepo, J. M., and Ullrich, P. A. (2023). A review of recent and emerging machine learning applications for climate variability and weather phenomena. *Artificial Intelligence for the Earth Systems*, 2(4), 220086.
- Sha, Y., Gagne II, D. J., West, G., and Stull, R. (2020). Deep-learning-based gridded downscaling of surface meteorological variables in complex terrain. Part I: Daily maximum and minimum 2-m temperature. *Journal of Applied Meteorology and Climatology*, 59(12), 2057-2073.
- Yu, S., Chakraborty, I., Anderson, G. J., Lucas, D. D., Lops, Y., and Galea, D. (2024). UFNNet: Joint U-Net and fully connected neural network to bias correct precipitation predictions from climate models. *Artificial Intelligence for the Earth Systems*.
- Wang, F., and Tian, D. (2022). On deep learning-based bias correction and downscaling of multiple climate models simulations. *Climate dynamics*, 59(11), 3451-3468.
- Ling, F., Li, Y., Luo, J. J., Zhong, X., and Wang, Z. (2022). Two deep learning-based bias-correction pathways improve summer precipitation prediction over China. *Environmental Research Letters*, 17(12), 124025.
- Zhang, Y., Chen, M., Han, L., Song, L., and Yang, L. (2022). Multi-element deep learning fusion correction method for numerical weather prediction. *Acta Meteorol. Sin*, 80, 153-167.
- Pan, B., Anderson, G. J., Goncalves, A., Lucas, D. D., Bonfils, C. J., Lee, J., Tian, Y., and Ma, H. Y. (2021). Learning to correct climate projection biases. *Journal of Advances in Modeling Earth Systems*, 13(10), e2021MS002509.
- Hess, P., Lange, S., Schötz, C., and Boers, N. (2023). Deep Learning for Bias-Correcting CMIP6-Class Earth System Models. *Earth's Future*, 11(10), e2023EF004002.
- Teutschbein, C., and Seibert, J. (2012). Bias correction of regional climate model simulations for hydrological climate-change impact studies: Review and evaluation of different methods. *Journal of hydrology*, 456, 12-29.
- Han, L., Chen, M., Chen, K., Chen, H., Zhang, Y., Lu, B., Song, L., and Qin, R. (2021). A deep learning method for bias correction of ECMWF 24–240 h forecasts. *Advances in Atmospheric Sciences*, 38(9), 1444-1459.
- Faijaroenmongkol, T., Sarinapakorn, K., and Vateekul, P. (2023). Sub-Seasonal Precipitation Bias-Correction in Thailand Using Attention U-Net With Seasonal and Meteorological Effects. *IEEE Access*, 11, 135463-135475.