

Unsupervised Image Classification Algorithms Applied to Fire-Prone Area Detection

I. Rahimi^{1,2}, L. Duarte^{1,2} and A. C. Teodoro^{1,2}

¹*Department of Geosciences, Environment and Spatial Planning Faculty of Sciences of the University of Porto, Portugal*

²*Institute of Earth Sciences, FCUP Pole, Porto, Portugal*

Keywords: Remote Sensing, Wildfire, Classification, Zagros Mountains.

Abstract: Remote sensing data has become critical in identifying fire-prone areas, providing essential insights through satellite imagery and various geospatial inputs. These data sources allow for real-time monitoring, mapping fire susceptibility, and assessing factors such as vegetation, fuel moisture, land use, and environmental conditions. Numerous supervised and unsupervised models combined with remote sensing data have shown great potential in predicting fire-prone regions, offering accurate and timely information for early warning systems and resource allocation. This study focuses on applying two unsupervised methods—PCA, and K-means—using inputs like Sentinel-2 imagery, elevation, and the Zagros Grass Index (ZGI) to identify fire-prone areas in the Kurdo-Zagrosian forests, an area increasingly vulnerable to wildfires. Among the two methods evaluated, PCA demonstrated superior performance in predicting fire-susceptible areas, accurately classifying 80% of the burned regions from 2021 to 2023 as moderate to high-risk zones.

1 INTRODUCTION


Forest fires' increasing frequency and intensity worldwide is an escalating concern, driven by natural and human-induced factors such as extreme weather conditions, shifting land use patterns, and rapid urban expansion (Zema, 2020; Bowman, 2017). These factors, especially under the growing influence of climate change, exacerbate the risk of wildfires. This significant loss of forest cover highlights the urgent need for effective wildfire monitoring and prevention strategies.


In recent years, integrating geospatial and remote sensing (RS) data/technologies has provided invaluable insights into wildfire risk factors (Teodoro, 2013). Through RS data, researchers can monitor and analyze variables such as land cover, temperature, and vegetation phenology (Duarte, 2018). This spatial data, combined with Geographic Information Systems (GIS), allows for the continuous monitoring of large areas and provides timely information about the likelihood of fire incidents (Mishra, 2024). RS data enables researchers to track environmental changes in real time and analyze


critical variables such as fuel moisture content, temperature trends, and human activity patterns, all of which contribute to the increased risk of fires.

A key application of this technological advancement is the development of Forest Fire Susceptibility Maps (FSMs). These maps are crucial for identifying areas at high risk of wildfires, enabling authorities to allocate resources efficiently and implement mitigation strategies in advance. By locating and assessing fire-prone regions, FSMs play a central role in reducing the vulnerability of ecosystems and communities while supporting informed ecological and urban planning decisions (Ghorbanzadeh, 2019). The effectiveness of FSMs has been significantly enhanced by integrating GIS, RS, and image classification algorithms, allowing researchers to predict fire susceptibility better and improve early warning systems (Rihan, 2023).

The growing popularity of image classification algorithms in fire susceptibility mapping is due to their ability to process large-scale, high-dimensional datasets and model complex non-linear relationships more effectively than traditional statistical methods (Kantarcioğlu, 2023). These models perform well at

^a <https://orcid.org/0009-0002-7411-8637>

^b <https://orcid.org/0000-0002-7537-6606>

^c <https://orcid.org/0000-0002-8043-6431>

integrating diverse data types, from climatic conditions and vegetation indices to human activity metrics, which allows for more precise fire risk assessments. In recent years, the development of FSM has been advanced through both single-model applications and ensemble methods, frequently showing improved performance over more conventional techniques (Piao, 2022; Saha, 2023). These advancements underscore FSM's potential to surpass traditional frameworks, particularly by leveraging methods that incorporate broader environmental and spatial analysis techniques.

Unsupervised classification methods, such as Principal Component Analysis (PCA) and K-means clustering, have proven valuable for extracting meaningful patterns from satellite imagery. PCA reduces large datasets into key components, capturing the most significant variations (Sharma, 2021), which is crucial for identifying fire susceptibility factors like vegetation, moisture, and land use patterns. K-Means, on the other hand, clusters satellite image data based on spectral similarity, enabling the identification of regions with similar environmental characteristics (Theodoridis, 2009). These methods are especially effective in handling large-scale, heterogeneous datasets from RS data (Celik, 2009).

The objective of this study is to apply two unsupervised image classification techniques—PCA and K-means—using inputs like Sentinel-2 imagery, elevation, and the Zagros Grass Index (ZGI) (Rahimi, 2024) to identify fire-prone areas in the Kurdo-Zagrosian forests. This region, vulnerable to increasing wildfire incidents, presents a complex environmental landscape where these methods can highlight patterns indicating higher fire susceptibility. By utilizing unsupervised algorithms, the study aims to provide a data-driven approach to wildfire susceptibility mapping, improving prevention and mitigation strategies in fire-prone regions.

2 METHODS

2.1 Study Area

The method is designed for areas with Semi-Arid (SA) and Semi-Mediterranean (SM) climates, exemplified by the forests of Marivan and Sarvabad in the Kurdistan Province, western Iran, bordering Iraq. These regions have experienced frequent fire incidents over the past decade. Geographically, the study area spans between longitudes 45°57'50" E and 46°46'41" E and latitudes 35°1'1" N and 35°49'51" N. Located in the northern Zagros mountains, Marivan

sits at an average elevation of 1,287 meters above sea level and features varied topography, from mountains to valleys, shaped by its cross-border proximity with Iraq (Rahimi, 2023).




Figure 1: Study area: Marivan and Sarvabad, Kurdistan province, Iran.

The climate in this region is SM, characterized by cold winters and hot summers. Annual rainfall averages 991 mm, with considerable variability (± 235 mm). Humidity averages around 54%, which is insufficient to sustain green pastures during summer, leaving behind dry grass that elevates fire risks (Rahimzadeh, 2008).

2.2 Data Collection and Preparation

Table 1 summarizes the data used in this study. Sentinel-2 satellite imagery and related products provided high-resolution multispectral data essential for environmental analysis and fire risk assessment.

Table 1: Data sources used.

Data Type, Projection System, Spatial Resolution (m)	Time Period	Data Source
Sentinel 2 (13 bands), UTM ¹ , 10	13/05/2020 and 15/09/2020	ESA ³
Sentinel Burned Area Product, UTM, 10	2021 -2023	
ZGI, UTM, 10	2020	(Rahimi, 2024)
DEM/SRTM ² , UTM, 10	-	USGS ⁴

1. Universal Transverse Mercator
2. Digital Elevation Model/Shuttle Radar Topography Mission
3. European Space Agency
4. United States Geological Survey

2.3 Methodology

The collected data layers were first stacked using Python 3.12, ensuring spatial alignment and consistency across all layers. To integrate Sentinel-2 and ZGI layers, the datasets were clipped to the boundaries of the study area using a defined polygon.

The clipped layers were then aligned to ensure consistent spatial resolution and extent. Finally, the processed layers were stacked into a multi-band raster, enabling the combination of spectral information and vegetation index data for subsequent fire susceptibility analysis. This approach ensured a precise and comprehensive dataset for modeling and assessment. These layers were integrated into a matrix consisting of all raster layers, which included Sentinel-2 bands (13 bands), elevation, and the ZGI. The resulting matrix had dimensions of 15×8020×6557, meaning that 15 different values represented each pixel in the study area. Figure 2 illustrates the methodology and process flow of this approach.




Figure 2: Methodological framework.

This study employed PCA and K-means to detect fire-prone areas effectively. To evaluate the effectiveness of these models, the Leave-One-Out (LOO) method was implemented, allowing for a thorough assessment of how different data layers contributed to fire susceptibility mapping. The validation of the results with post-2020 data (from

2021–2023) demonstrated the reliability of these methods for accurately predicting fire-prone areas over time, showcasing their potential for long-term wildfire risk assessment and management.

K-means and PCA were chosen as they are common unsupervised methods suitable for studies with limited field data. Unlike supervised learning approaches, which require extensive datasets and computational resources, these methods provide a practical and efficient framework for fire susceptibility classification. The components were retained to classify fire susceptibility into high, average, and low categories. This decision reflects the study area's vegetation and land cover characteristics alongside the limited field data available. This approach balances capturing sufficient data variance with the need for interpretability, ensuring robust and practical classification.

To validate the results, the post-2020 burned area was used. They derived from sentinel 2 by applying the Normalize Burned Ratio (NBR) on data from 2021 to 2023 (Giddey, 2021). The NBR index proved to be a valuable tool for identifying burned areas by leveraging spectral differences between vegetation and charred surfaces. This methodology ensured reliable validation of the predictive models and reinforced the importance of integrating remote sensing indices into fire susceptibility research.

3 RESULTS

Figure 3 shows the results of applying PCA and K-means. The results revealed varying levels of effectiveness (High, Average, and Low) in predicting fire-prone areas in the study region using 2020 data. The black polygons are the burned areas calculated from Sentinel 2 data from 2021 to 2023, using the NBR index.

These patterns emphasize the spatial variability of fire-prone areas, which is influenced by factors such as vegetation type, fuel density, and topographic conditions, as suggested by earlier studies (Rahimi et al., 2024). The classification levels in this study were established based on the number of components selected for each of the unsupervised methods, PCA and K-means, with three components defined for optimal model performance in both cases. The trade-off between preserving critical information and minimizing computational complexity informed this selection process for the number of components. By reducing the dimensionality of the data, both PCA and K-means enhanced interpretability while maintaining the integrity of key predictive features (Giddey, 2021).




Figure 3: Maps generated by a) PCA and b) K-means according to data in 2020. The black polygons demonstrate the burned areas from 2021 to 2023.

The visual representation in Figure 3 highlights distinct patterns of fire susceptibility zones, with clear differentiation between regions of varying risk levels.

Among the methods, PCA produced the most accurate and reliable results, as shown in Figure 3a,

compared to K-means (Figure 3b). The red areas show high-fire susceptible areas, while the yellow and green colors show the average and low-fire susceptible areas. The stark contrast between the red zones and the remaining areas underscores the ability of PCA to highlight regions requiring urgent attention. This distinction aligns with practical applications in fire management, where prioritizing high-risk zones can significantly enhance resource allocation and mitigation efforts.

The validation process compared high-susceptibility areas identified by the models with actual burned areas in subsequent years. PCA exhibited a higher overlap between its predicted high fire-susceptible zones and the areas affected by post-2020 wildfires, with approximately 41%. In contrast, this value was 31.3% for the K-means (Figure 4). This significant difference in overlap suggests that PCA's ability to identify subtle patterns in the data gives it a clear edge over K-means. The higher accuracy of PCA aligns with its proven effectiveness in environmental modeling and its capacity to extract critical components from high-dimensional datasets. Regarding High and Average fire-prone areas together, both methods offered almost the same results. This indicates that PCA outperformed the K-means models in accurately detecting fire-prone regions, highlighting its robustness in mapping fire susceptibility. However, the similar performance of the two methods in identifying combined High and Average susceptibility areas suggests that K-means may still hold utility in broader-scale applications where granular accuracy is less critical. This underscores the need for selecting methods based on specific use-case requirements.




Figure 4: Detection of fire-prone areas for the post-2020 years by 2020's data using PCA and K-means.

Moreover, the analysis indicated that certain Sentinel-2 spectral bands could be excluded without substantially reducing the model's accuracy. This

finding highlights the efficiency of PCA in dimensionality reduction, where less critical spectral bands are discarded while retaining the most informative features. Such optimization reduces computational demands and facilitates faster data processing, which is crucial for large-scale studies (Kantarcioglu, 2023). This suggests that PCA effectively reduced dimensionality, preserving only the most informative features while maintaining high predictive accuracy. The ability to maintain predictive accuracy while reducing data complexity makes PCA an attractive option for fire susceptibility mapping, particularly in resource-constrained settings where computational efficiency is essential. The fact that the PCA model consistently provided reliable predictions across multiple years demonstrates its potential for long-term application in fire susceptibility mapping.

This consistency underscores PCA's robustness and adaptability, especially in evolving environmental conditions and fire patterns. Its reliability across diverse temporal datasets reaffirms its suitability for integration into long-term fire management frameworks.

4 CONCLUSIONS

In conclusion, this study highlights the effectiveness of using RS data combined with unsupervised image classification techniques to identify fire-prone areas in the Kurdo-Zagrosian forests. Among the two methods tested, PCA performed better in accurately predicting fire-susceptible zones, with 80% of the burned areas from 2021 to 2023 correctly classified as moderate to high-risk. Its ability to reduce dimensionality while preserving critical information, especially when excluding less relevant Sentinel-2 bands, further solidifies PCA as a robust tool for long-term fire risk mapping. In contrast, K-means showed moderate success, identifying around 50% of the burned areas. These findings emphasize the potential of PCA for improving fire management strategies, while K-means may require further refinement to achieve similar predictive accuracy.

In the future, more unsupervised classification methods will be compared to further evaluate their effectiveness and explore potential improvements in fire risk prediction.

ACKNOWLEDGMENTS

The work is funded by national funds through FCT – Fundação para a Ciência e Tecnologia, I.P., in the framework of the UIDB/04683/2020 (<https://doi.org/10.54499/UIDB/04683/2020>) and UIDP/04683/2020 633 (<https://doi.org/10.54499/UIDP/04683/2020>) – Instituto de Ciências da Terra programs.

REFERENCES

- Bowman, M.J.S.D.; Moreira-Munoz, A.; Kolden, C.A.; Chavez, R.O.; Munoz, A.A.; Salinas, F.; Gonzalez-Reyes, A.; Rocco, R.; de la Barrera, F.; Williamson, G.J.; et al. Human- Environmental Drivers and Impacts of the Globally Extreme 2017 Chilean Fires. *Ambio* 2018, doi:10.1007/s13280-018-1084-1
- Celik, N. T. (2009). Unsupervised Change Detection in Satellite Images Using Principal Component Analysis and K-Means Clustering. *IEEE Geoscience and Remote Sensing Letters*, 6(4), 772–776. <https://doi.org/10.1109/lgrs.2009.2025059>
- Duarte, L., Teodoro, A., Monteiro, A.T., Cunha, C., & Gonçalves, H., "PhenoMetrics: An open source software application to assess vegetation phenology metrics," *Computers and Electronics in Agriculture* 148, 82–94 (2018); <https://doi.org/10.1016/j.compag.2018.03.007>.
- ESA European Space Agency (ESA), "Copernicus Open Access Hub, Sentinel-2 Level-2A products," <https://scihub.copernicus.eu>.
- Ghorbanzadeh, O.; Valizadeh, K.K.; Blaschke, T.; Aryal, J.; Naboureh, A.; Einali, J.; Bian, J. Spatial prediction of wildfire susceptibility using field survey GPS data and machine learning approaches. *Fire* 2019, 2(3), 43. <https://doi.org/10.3390/fire2030043>.
- Giddey, B. L., Baard, J. A., & Kraaij, T. (2021). Verification of the differenced Normalised Burn Ratio (dNBR) as an index of fire severity in Afrotropical Forest. *South African Journal of Botany*, 146, 348–353. <https://doi.org/10.1016/j.sajb.2021.11.005>
- Kantarcioglu, O.; Schindler, K.; Kocaman, S. Forest fire susceptibility assessment with machine learning methods in north-east Türkiye. *Int. Arch. Photogramm. Remote Sens. Spatial Inf. Sci.* 2023, XLVIII-M-1-2023, 161–167. <https://doi.org/10.5194/isprs-archives-xxviii-m-1-2023-161-2023>.
- Mishra, M.; Guria, R.; Baraj, B.; Nanda, A.P.; Santos, C.A.G.; Da Silva, R.M.; Laksono, F.A.T. Spatial analysis and machine learning prediction of forest fire susceptibility: A comprehensive approach for effective management and mitigation. *Sci. Total Environ.* 2024, 926, 171713. <https://doi.org/10.1016/j.scitotenv.2024.171713>.
- Piao, Y.; Lee, D.; Park, S.; Kim, H.G.; Jin, Y. Forest fire susceptibility assessment using Google Earth Engine in Gangwon-do, Republic of Korea. *Geomatics Nat.*

- Hazards Risk* 2022, 13, 432–450. <https://doi.org/10.1080/19475705.2022.2030808>.
- Rahimi, I., Duarte, L., & Teodoro, A. C. (2024). Zagros Grass Index—A new vegetation index to enhance fire fuel mapping: A case study in the Zagros Mountains. *Sustainability*, 16(10), 3900. <https://doi.org/10.3390/su16103900>
- Rahimzadeh Bajgiran, P., Darvishsefat, A., Khalili, A., & Makhdoum, M., (2008). "Using AVHRR-based vegetation indices for drought monitoring in the Northwest of Iran," *Journal of Arid Environments* 72, 1086–1096
- Rihan, M.; Bindajam, A.A.; Talukdar, S.; Shahfahad; Naikoo, M.W.; Mallick, J.; Rahman, A. Forest fire susceptibility mapping with sensitivity and uncertainty analysis using machine learning and deep learning algorithms. *Advances in Space Research* 2023, 72(2), 426–443. <https://doi.org/10.1016/j.asr.2023.03.026>.
- Saha, S.; Bera, B.; Shit, P.K.; Bhattacharjee, S.; Sengupta, N. Prediction of forest fire susceptibility applying machine and deep learning algorithms for conservation priorities of forest resources. *Remote Sensing Applications: Society and Environment* 2023, 29, 100917. <https://doi.org/10.1016/j.rsase.2022.100917>.
- Sharma, D., Sharma, T., & Singhai, J. (2021). Extraction of Water Bodies from Visible Color Satellite Images Using PCA Feature Map. *2021 IEEE International India Geoscience and Remote Sensing Symposium (InGARSS)*, 1–4. <https://doi.org/10.1109/ingarss51564.2021.9791857>
- Teodoro, A.C., Duarte, L., (2013). Forest fire risk maps: a GIS open source application a case study in Norwest of Portugal, *International Journal of Geographical Information Science* 27(4), DOI:10.1080/13658816.2012.721554.
- Theodoridis, S., & Koutroumbas, K. (2009). Clustering Algorithms III: Schemes based on Function Optimization. In *Elsevier eBooks* (pp. 701–763). <https://doi.org/10.1016/b978-1-59749-272-0.50016-5>
- USGS EROS Archive - Digital Elevation - Shuttle Radar Topography Mission (SRTM), <https://www.usgs.gov/centers/eros/science/usgs-eros-archive-digital-elevation-shuttle-radar-topography-mission-srtm-1>.
- Zema, D.A.; Nunes, J.P.; Lucas-Borja, M.E, (2020). Improvement of Seasonal Runoff and Soil Loss Predictions by the MMF (Morgan-Morgan-Finney) Model after Wildfire and Soil Treatment in Mediterranean *Forest Ecosystems. Catena*, 188, 104415, doi: 10.1016/j.catena.2019.104415