Predicting Photovoltaic Power Output Using LSTM: A Comparative Study Using both Historical and Climate Data

Fereshteh Jafari^{1,2}¹, Joseph Moerschell¹¹, and Kaspar Riesen²

¹Institute of System Engineering, University of Applied Sciences and Arts Western Switzerland, Sion, Switzerland ²Institute of Computer Science, University of Bern, CH-3012 Bern, Switzerland

- Keywords: LSTM, Meteorological Data Integration, Photovoltaic Power Prediction, Prediction Horizon Assessment, Temporal Sliding Window Analysis.
- Abstract: Accurate photovoltaic (PV) power output prediction is important for efficient energy management in solar power systems. This study explores the benefits and limitations of Long Short-Term Memory (LSTM) networks in predicting PV power using three distinct approaches, namely using historical PV power data, climate data, and a combination of both, all with timestamps. The performance of these methods is evaluated across different prediction horizons of 10, 30, and 50 minutes ahead. Additionally, the impact of the sliding window size, representing the amount of past data used for training, is analyzed. The models are trained and tested on a dataset collected over three months from a rooftop PV system in Sion, Switzerland, with a maximum power of 22.2 kW. The Root Mean Square Error as well as the *R*² metrics are provided to assess the accuracy of each method. The results demonstrate that both the choice of the actual input data and the sliding window size significantly influence the prediction accuracy. In particular, the results presented here show the potential of combining different data sources to improve the accuracy of PV power prediction using LSTM models.

1 INTRODUCTION

1.1 Problem Definition

Solar power stands out due to its cleanliness, wide availability, and scalability. However, photovoltaic (PV) power output is inherently variable across multiple timescales due to factors such as geographical location, as well as the size, orientation and technology of the panels. Moreover, environmental conditions like solar irradiance, temperature, cloud cover, wind, and humidity crucially influence PV power output (Pelland et al., 2013), (Pedro and Coimbra, 2012). These characteristics pose significant challenges for grid stability and energy management (Chen and Chang, 2021). Moreover, according to the World Energy Outlook 2024 by the International Energy Agency, global energy demand is expected to increase by approximately 16% by 2050 under the Stated Policies Scenario (International Energy Agency, 2024). However, if energy efficiency measures are not enhanced, this demand could rise by about 22% over the same period. Accurate short-term PV power forecasts are thus critical not only for operational efficiency but also for effective energy storage, load control, and system reliability contributing to the broader goals of energy conservation, sustainability, and climate change mitigation (Park et al., 2021). Consequently, the development of robust predictive models for PV power generation throughout the day is important to address these challenges.

1.2 State of the Art

Prediction methods for PV power output span from traditional approaches like persistence, physical, and statistical models, which use both current PV data and weather data, to more sophisticated models that employ artificial intelligence (AI). Persistence models commonly rely on present data for immediate predictions (Wang et al., 2021), (Dash et al., 2021), while physical models incorporate weather variables (Lima et al., 2016). Statistical models (Wang et al., 2022), such as ARMA (Benmouiza and Cheknane, 2016) and ARIMA (Reikard, 2009), struggle with nonstationary and nonlinear data, limitations that more ad-

In Proceedings of the 14th International Conference on Pattern Recognition Applications and Methods (ICPRAM 2025), pages 733-740 ISBN: 978-989-758-730-6; ISSN: 2184-4313

^a https://orcid.org/0000-0001-6049-6536

^b https://orcid.org/0000-0003-3072-6075

^c https://orcid.org/0000-0002-9145-3157

Jafari, F., Moerschell, J. and Riesen, K.

Predicting Photovoltaic Power Output Using LSTM: A Comparative Study Using both Historical and Climate Data. DOI: 10.5220/001325800003905

Copyright © 2025 by Paper published under CC license (CC BY-NC-ND 4.0)

vanced AI models aim to overcome (Kumbhar et al., 2021).

AI technologies, including machine learning (ML), are important to improve building energy management systems as well as energy efficiency. ML can broadly be divided into several branches, including supervised and unsupervised learning, reinforcement learning, and deep learning (Mellit et al., 2020). Among these, supervised learning is the most prevalent approach in ML, where models are trained on labeled datasets that include both predictors and predictands (Markovics and Mayer, 2022). The overall goal of supervised learning is to identify the optimal functional relationship between input variables and outputs, enabling accurate predictions. Supervised ML problems can generally be divided into two categories, viz. classification, which deals with categorical outputs, and regression, which addresses continuous outputs. PV power prediction, specifically, is categorized as a regression problem due to its focus on predicting continuous power output values (Wang et al., 2023).

Common ML models used for PV power predictions include artificial neural networks (ANN), support vector regression (SVR), k-nearest neighbors (KNN) regression, as well as linear regression (Mellit et al., 2020). While ANN models can capture complex nonlinear relationships by utilizing large datasets, they may often not fully address the dynamic characteristics of PV power generation, which is influenced by various temporal factors (Meenal and Selvakumar, 2018). As a result, alternative methods such as SVR and KNN also attempt to model these nonlinear relationships, but they can struggle with the inherent time-dependent nature of the data (Wang et al., 2017), (Liu et al., 2018), and (Mohammadi et al., 2015). Deep learning models like recurrent neural networks (RNNs) and long-short term memory (LSTM) effectively address these challenges by capturing sequential dependencies in data. While RNNs excel in time-series prediction, they struggle with issues like vanishing gradients, LSTMs may overcome these problems, thereby enhancing the accuracy of PV power prediction (Wang et al., 2020), (Wang et al., 2019).

The superiority of LSTM models over other ML methods for the prediction of PV power has been empirically proven in several studies. In (Gao et al., 2019), for instance, two distinct LSTM-based prediction models tailored for ideal and non-ideal weather conditions are introduced. For ideal weather, the model leverages numerical weather prediction data with seasonal adjustments, achieving an RMSE accuracy of 4.62%. For non-ideal weather conditions,

the model integrates a discrete grey model to predict daily total power, enhancing the LSTM's accuracy in scenarios such as rainy, cloudy, and overcast days. In a second study, a one-hour ahead prediction of PV power using historical PV power data only is proposed (Abdel-Nasser and Mahmoud, 2019). The authors compare the performance of the LSTM model with three PV prediction methods for stationary time series. Then, in (Hu et al., 2024) historical PV power data and climate data is used separately to predict the PV power. Results show that the use of climate data leads in general to better results with a smaller RMSE and the highest R^2 values. Finally, the importance of selecting predictors based on model architecture in solar energy forecasting is highlighted in (Ciobanu et al., 2024). A dual-view LSTM model, using both historical and future weather data, outperforms a single-view model that only uses historical PV production and past weather data. Humidity consistently emerges as an important predictor across both models, underscoring the value of future weather data in enhancing predictive accuracy.

1.3 Contribution

While several studies demonstrate the potential of LSTMs in the domain of PV power prediction, they often focus on single data sources or overlook the influence of input data configurations on prediction accuracy (e.g., the sliding window size). The present paper provides a time-series prediction that seeks to bridge these gaps. First, we thoroughly evaluate the performance of LSTM models trained with different data sources-past PV power, climate data, and their combination-across multiple prediction horizons and discuss on how these inputs effect the predictions. Second, the study also examines the effect of varying sliding window sizes on the prediction accuracy. That is, we provide a nuanced understanding of how temporal context influences predictive performance by providing empirical evidence using realworld PV power data (adding practical relevance to the findings of this paper).

1.4 Paper Organization

The remainder of this paper is organized as follows. Section 2 outlines the methodology of the work, including data type and preparation. Section 3 presents the predictive model architecture. In Section 4 we show and discuss the experimental results. Finally, Section 5 concludes the paper with key insights.

2 METHODOLOGY

2.1 Long-Short Term Memory Network (LSTM)

One of the major advantages of RNNs is their capacity to incorporate contextual information when mapping between input and output sequences, making them particularly suitable for sequence prediction tasks. However, traditional RNNs face a limitation in effectively accessing long-term context, as the influence of a given input decays or amplifies exponentially with each recurrence, leading to what is known as the vanishing or exploding gradient problem (Graves, 2012). Figure 1 shows a simple RNN containing a single, self connected hidden layer.



Figure 1: A simple RNN containing a single, self connected hidden layer (Graves, 2012).

LSTM networks (Hochreiter and Schmidhuber, 1997) are a type of RNN designed to handle longterm dependencies using memory blocks instead of traditional neurons. These blocks incorporate forget, input, and output gates to manage the flow of information, addressing gradient decay and explosion issues. This makes LSTMs highly effective for sequence prediction tasks, such as PV power prediction, where capturing long-term patterns is crucial.

Figure 2(a) illustrates a single-cell LSTM memory block, showing how an LSTM network structure differs from a standard RNN by substituting the summation units in the hidden layer with memory blocks, as seen in Figure 2(b). Although ordinary summation units can be combined with LSTM blocks, this is generally unnecessary. Additionally, LSTM networks utilize the same output layers as those in standard RNNs (Graves, 2012).

Training of LSTM networks is conducted through backpropagation through time, which allows the network to learn from sequential data by minimizing the prediction error over multiple time steps.

2.2 Description of the PV Power System

In this section we briefly describe the PV power generation system used in this study, located at the authors' institution rooftop in Sion, Switzerland. Part of the setup is shown in Figure 3. The PV system has an installed maximum power of 22.2 kW and comprises multiple solar panels positioned to capture maximum sunlight throughout the day. The PV array consists of 60 individual solar panels, each with a rated power of 370 W. The PV modules use mono-crystalline silicon technology. There are two AMPT converters feeding a DC bus operating at 760 V. This DC bus connect different loads and storage batteries. Table 1 summarizes the specifications of an individual PV panel.

2.3 Data Collection and Preprocessing

This study uses real-time climate data from MeteoSwiss, including solar irradiance, temperature, wind speed, humidity, and sunshine duration, recorded at 10-minute intervals in Sion, Switzerland, from July 5 to September 30, 2024. The PV power production data, unique to this research, is obtained from a platform developed at GridLab, HES-SO Valais, and synchronized with the climate data for time-series analysis. Comprehensive data preprocessing is manually conducted, including checks for missing values, synchronization of climate data with PV power output, and normalization. This study also considers timestamps as features to train the model.

2.4 Sliding Window and Prediction Horizons

Major goal of this paper is to predict the PV power output at different prediction horizons, viz. 10, 30, and 50 minutes into the future. These prediction horizons are selected to provide short-term forecasts, which are valuable for real-time grid management and energy planning. To achieve these predictions, we utilize a sliding window approach in the LSTM model, where the length of historical data used for prediction (i.e., the sliding window) varies between 60 and 20 minutes. The chosen window size potentially influences the model's accuracy, as it dictates the amount of past information utilized to make each prediction. Moreover, it is highly dependent on the type of data actually employed to predict the PV power. In this study, we consider three different types of data:

- Climate data
- Historical PV power data
- Combination of climate data and historical PV power data



Figure 2: An LSTM network, (a) memory block with one cell, (b) with four input units, a hidden layer of two single-cell LSTM memory blocks and five output units (Graves, 2012).

Table 1: Specifications of an individual PV panel used in this study.

Modules	$P_{max}(W)$	$V_{oc}(V)$	$I_{sc}(A)$	$V_{mpp}(V)$	$I_{mpp}\left(A ight)$
LG370Q1C-V5	370	42.8	10.82	37.0	10.01



Figure 3: The studied system in Sion, Switzerland.

2.5 Evaluation Metrics

The Root Mean Square Error (RMSE) serves as a widely recognized metric for regression model evaluation. The RMSE is, for instance, employed as a standard statistical metric to assess model performance in fields such as meteorology, air quality, and climate research (Hodson, 2022). For a sample of *n* observations *y* (y_i , i = 1, 2, ..., n), and *n* corresponding model predictions \hat{y} , the RMSE is defined as

RMSE =
$$\sqrt{\frac{1}{n} \sum_{i=1}^{n} (y_i - \hat{y}_i)^2}$$
 (1)

For PV power prediction, RMSE provides a clear indication of the overall model accuracy (the smaller the RMSE value, the better the model).

The coefficient of determination R^2 is another widely used metric in regression model evaluation due to its intuitive interpretability (despite not always being the primary criterion for model selection). Formally, the R^2 coefficient is defined as follows

$$R^{2} = 1 - \frac{\sum_{i=1}^{n} (y_{i} - \hat{y}_{i})^{2}}{\sum_{i=1}^{n} (y_{i} - \bar{y})^{2}},$$
(2)

where y_i represents the observed values, \hat{y}_i the predicted values, \bar{y} is the mean of the observed values, and *n* is the number of observations.

The R^2 coefficient indicates the proportion of variance in the observed data that is predictable from the independent variables, making it valuable in understanding the model's real-world applicability (Gao, 2024). An R^2 value closer to 1 suggests a model that better captures the underlying patterns of the data.

3 PROPOSED WORK: PREDICTIVE MODEL

This section describes the modeling approach for predicting PV power output using various input data sources and the LSTM model's structure. It outlines three prediction models designed to assess the impact of different data sources on prediction accuracy, as discussed in Section 2.4. Additionally, it details the LSTM model architecture and training procedures specific to each model.

Figure 4 presents an overview of the entire modeling process in a flowchart format. The modeling process is split in two major phases, the data preprocessing and the PV power prediction. The PV power prediction phase is thoroughly explained in the next subsection.

The preprocessing consists of two steps. The first step involves checking for missing entries in both the PV power output and climate data to identify gaps and address inconsistencies. In the climate dataset, recorded at 10-minute intervals, missing values are rare and only occur between two recorded points. These missing entries are imputed with the mean of the preceding and succeeding values, ensuring a smooth temporal continuity in the climate data. PV power output data, is originally recorded every minute. Thus, the second step consist of synchronization with the climate data to maintain consistent intervals across datasets. To match the 10-minute intervals of the climate data, the average PV power is calculated over each 10-minute span, starting from the same timestamp as the climate dataset.



Figure 4: Flowchart of the LSTM model.

3.1 Prediction Models

Three models are developed to predict PV power output for 10, 30, and 50-minute horizons using different input data to assess their impact on prediction accuracy.

3.1.1 Model 1: PV Power-Only Prediction

Model 1 uses historical PV power data as the sole input for predicting future PV power output. By leveraging the temporal patterns in past PV power values, this model aims to understand the degree to which prior PV output data alone can predict shortterm power generation. This model assumes that PV power generation is somewhat self-correlated, capturing daily cycles and other temporal patterns without the influence of weather variables.

3.1.2 Model 2: Climate Data-Only Prediction

Model 2 is based exclusively on climate data, which includes features such as solar irradiance, temperature, wind speed, humidity, and sunshine duration. These features are collected from the MeteoSwiss database at a 10-minute interval and synchronized with the actual PV output data. By using climate data only, this model evaluates the predictive power of environmental conditions on PV generation, independent of past PV power values. Climate data is often highly correlated with PV output, making it a key predictor of solar energy production.

According to the correlation analysis provided in Figure 5, we can see that irradiance and sunshine duration are the most effective inputs in this endeavor with correlations of about 0.95 and 0.80, respectively.



Figure 5: Correlation map between climate data and PV power production.

3.1.3 Model 3: Combined Data Prediction

Model 3 combines both past PV power and climate data to improve the prediction accuracy by integrating both temporal patterns and environmental influences. This approach leverages the complementary information between historical PV production data and weather conditions, aiming to capture complex relationships that could affect the PV output.

3.2 LSTM Model Architecture

Each of the three models shares a similar LSTM architecture, differing only in the input layer based on the data configuration. The architecture includes four LSTM layers with 50 hidden units, for learning temporal dependencies and capturing patterns. A fully connected layer translates LSTM features to the regression output layer, which produces continuous PV power predictions.

3.3 Training Procedure

Each model is trained using backpropagation through time to optimize the weights, with the goal of minimizing the prediction error. The complete dataset is divided into training and testing subsets, with 90% of the data used for training and 10% for testing. To train the LSTM model, the Adam optimizer is used (known for its efficiency and adaptability, especially in handling noisy data and sparse gradients (Diederik, 2014)). The model is trained for a maximum of 200 epochs with an initial learning rate of 0.01 to balance between convergence speed and stability during the training.

4 RESULTS AND DISCUSSION

This section presents an analysis of the performance of the three LSTM models under the discussed configurations.

4.1 Estimation Model

The performance of Model 2, which predicts PV power using only current climate data (horizon = 0), is analyzed. Figure 6 shows estimated versus actual PV power, with stem plots and RMSE values quantifying performance. The model accurately distinguishes between day and night, with no errors at night, and follows daytime PV power closely, maintaining a mean deviation below 1.5 kW in over 90% of cases. This proof of concept demonstrates that estimating current PV power using weather data is feasible. Future experiments will explore prediction horizons of 10, 30, and 50 minutes.



Figure 6: Results of the LSTM model to estimate the PV power with climate data and time stamps tested for 9 days with the stem plot and an RMSE of about 1kW.

4.2 Comparative Analysis of Prediction Models

The effectiveness of each prediction model is evaluated by comparing actual versus predicted PV power values for models 1, 2, and 3, as shown in Figures 7, 8, and 9. All models use the past 20 minutes of PV power data, climate data, their combination, and timestamps to predict PV power for a 10-minute horizon. While all models perform well, the model combining historical data with climate data generally yields the best results in qualitative analysis. The three LSTM models are also used to predict PV power for 30 and 50 minutes ahead. The RMSE and R^2 values are shown in Figure 10 and 11, respectively. The results indicate that, across all prediction horizons, utilizing a combination of historical PV power data and climate data consistently leads to more accurate PV power predictions. That is, this combined approach appears to enhance the model's ability to capture both short-term trends from recent PV output and underlying patterns influenced by climatic conditions, resulting in generally improved prediction accuracy.



Figure 7: A LSTM model to predict the PV power with historical data and timestamps with prediction horizon of 10 minutes.



Figure 8: A LSTM model to predict the PV power with climate data and timestamps with prediction horizon of 10 minutes.



Figure 9: A LSTM model to predict the PV power with historical, climate data and timestamps with prediction horizon of 10 minutes.

	Window sizes	Combination of both data							
		Predict 10 min ahead		Predict 30 min ahead		Predict 50 min ahead			
		RMSE (%)	R^2	RMSE (%)	R^2	RMSE (%)	R^2		
	20 min	4.39	0.93	6.25	0.86	6.67	0.84		
	60 min	4.50	0.93	6.52	0.85	7.19	0.82		

Table 2: Performance metrics for PV power prediction using a combination of historical and climate data across two sliding window sizes are presented, with the best results highlighted in bold.



Figure 10: RMSE values for different LSTM models to predict the PV power.



Figure 11: R^2 values for different LSTM models to predict the PV power, starting from 0.6.

4.3 Impact of Sliding Window Size

An additional experiment evaluates the impact of a 60 minutes sliding window size compared to the 20minute window used previously. The results of this experiment are shown in Table 2, where we present the RMSE and R^2 values for both window sizes, namely 20 and 60 minutes (for the sake of simplicity, we only show the results of combining both inputs, historical data, and climate data - similar effects are to be expected for the individual data sets). We find that for small prediction horizon (20-minute), the results are slightly better. However, for longer prediction horizon (60-minute), the higher RMSE and lower R^2 values for the 60-minute window size are clear. The nonlinear and highly oscillatory nature of climate data apparently results in suboptimal model training at long prediction horizons when a larger moving window is used. Perhaps when a shadow passes, it is more important to detect the start of the shadow than to remember what happened in the last 60 minutes (and the beginning of the shadow can best be recognized based on the current and past PV power generation values). Overall, these results suggest that a smaller sliding window can better capture relevant short-term dependencies when predicting PV power in longer prediction horizons.

5 CONCLUSIONS

This paper investigates the accuracy of PV power prediction using LSTM models trained with different datasets, namely historical PV power data, climate data, a combination of both all with timestamps. Prediction horizons of 10, 30, and 50 minutes are explored to assess model performance under varying conditions, and two different sliding window sizes are considered to show the effect of time and data type in prediction. Results indicate that RMSE values increase as the prediction horizon extends. The combined dataset of PV power and climate data with timestamps consistently yields lower RMSE values compared to using either dataset individually, indicating that integrating diverse data sources enhances prediction accuracy. The results also underscore the importance of selecting an appropriate sliding window size. These findings contribute valuable insights into optimizing LSTM models for short-term PV power prediction, with practical implications for real-world solar energy prediction.

ACKNOWLEDGEMENTS

This research is funded by the Swiss National Science Foundation (SNSF) under grant no. 10CE-1_216893 in cooperation with CETPartnership, under Joint Call 2022 for research proposals, Increasing control and efficiency in regional energy systems using quantum sensors and machine learning – QuantumIRES, UE-FISCDI PNCDI IV.

REFERENCES

- Abdel-Nasser, M. and Mahmoud, K. (2019). Accurate photovoltaic power forecasting models using deep LSTM-RNN. *Neural Comput. Appl.*, 31(7):2727–2740.
- Benmouiza, K. and Cheknane, A. (2016). Small-scale solar radiation forecasting using arma and nonlinear autoregressive neural network models. *Theoretical and Applied Climatology*, 124:945–958.
- Chen, H. and Chang, X. (2021). Photovoltaic power prediction of lstm model based on pearson feature selection. *Energy Reports*, 7:1047–1054.
- Ciobanu, A., Bacanin, N., Sanders, T., and Stoean, C. (2024). Assessing predictor influence in lstm models for enhanced solar energy forecasting. In Proceedings of the 26th International Symposium on Symbolic and Numeric Algorithms for Scientific Computing (SYNASC).
- Dash, D. R., Dash, P., and Bisoi, R. (2021). Short term solar power forecasting using hybrid minimum variance expanded rvfln and sine-cosine levy flight pso algorithm. *Renewable Energy*, 174:513–537.
- Diederik, P. K. (2014). Adam: A method for stochastic optimization. (*No Title*).
- Gao, J. (2024). R-squared (r2)-how much variation is explained? Research Methods in Medicine & Health Sciences, 5(4):104–109.
- Gao, M., Li, J., Hong, F., and Long, D. (2019). Day-ahead power forecasting in a large-scale photovoltaic plant based on weather classification using lstm. *Energy*, 187:115838.
- Graves, A. (2012). Long short-term memory. Supervised sequence labelling with recurrent neural networks, pages 37–45.
- Hochreiter, S. and Schmidhuber, J. (1997). Long short-term memory. *Neural Comput.*, 9(8):1735–1780.
- Hodson, T. O. (2022). Root mean square error (rmse) or mean absolute error (mae): When to use them or not. *Geoscientific Model Development Discussions*, 2022:1–10.
- Hu, Z., Gao, Y., Ji, S., Mae, M., and Imaizumi, T. (2024). Improved multistep ahead photovoltaic power prediction model based on lstm and self-attention with weather forecast data. *Applied Energy*, 359:122709.
- International Energy Agency (2024). World energy outlook 2024.
- Kumbhar, A., Dhawale, P. G., Kumbhar, S., Patil, U., and Magdum, P. (2021). A comprehensive review: Machine learning and its application in integrated power system. *Energy Reports*, 7:5467–5474.
- Lima, F. J., Martins, F. R., Pereira, E. B., Lorenz, E., and Heinemann, D. (2016). Forecast for surface solar irradiance at the brazilian northeastern region using nwp model and artificial neural networks. *Renewable Energy*, 87:807–818.
- Liu, L., Zhao, Y., Chang, D., Xie, J., Ma, Z., Sun, Q., Yin, H., and Wennersten, R. (2018). Prediction of shortterm pv power output and uncertainty analysis. *Applied energy*, 228:700–711.
- Markovics, D. and Mayer, M. J. (2022). Comparison of machine learning methods for photovoltaic

power forecasting based on numerical weather prediction. *Renewable and Sustainable Energy Reviews*, 161:112364.

- Meenal, R. and Selvakumar, A. I. (2018). Assessment of svm, empirical and ann based solar radiation prediction models with most influencing input parameters. *Renewable Energy*, 121:324–343.
- Mellit, A., Massi Pavan, A., Ogliari, E., Leva, S., and Lughi, V. (2020). Advanced methods for photovoltaic output power forecasting: A review. *Applied Sciences*, 10(2):487.
- Mohammadi, K., Shamshirband, S., Anisi, M. H., Alam, K. A., and Petković, D. (2015). Support vector regression based prediction of global solar radiation on a horizontal surface. *Energy Conversion and Management*, 91:433–441.
- Park, M. K., Lee, J. M., Kang, W. H., Choi, J. M., and Lee, K. H. (2021). Predictive model for pv power generation using rnn (lstm). *Journal of Mechanical Science* and Technology, 35(2):795–803.
- Pedro, H. T. and Coimbra, C. F. (2012). Assessment of forecasting techniques for solar power production with no exogenous inputs. *Solar Energy*, 86(7):2017–2028.
- Pelland, S., Remund, J., Kleissl, J., Oozeki, T., and De Brabandere, K. (2013). Photovoltaic and solar forecasting: state of the art. *IEA PVPS Task*, 14(355):1–36.
- Reikard, G. (2009). Predicting solar radiation at high resolutions: A comparison of time series forecasts. *Solar energy*, 83(3):342–349.
- Wang, F., Lu, X., Mei, S., Su, Y., Zhen, Z., Zou, Z., Zhang, X., Yin, R., Duić, N., Shafie-khah, M., et al. (2022). A satellite image data based ultra-short-term solar pv power forecasting method considering cloud information from neighboring plant. *Energy*, 238:121946.
- Wang, F., Xuan, Z., Zhen, Z., Li, K., Wang, T., and Shi, M. (2020). A day-ahead pv power forecasting method based on lstm-rnn model and time correlation modification under partial daily pattern prediction framework. *Energy Conversion and Management*, 212:112766.
- Wang, F., Zhen, Z., Wang, B., and Mi, Z. (2017). Comparative study on knn and svm based weather classification models for day ahead short term solar pv power forecasting. *Applied Sciences*, 8(1):28.
- Wang, H., Lei, Z., Zhang, X., Zhou, B., and Peng, J. (2019). A review of deep learning for renewable energy forecasting. *Energy Conversion and Management*, 198:111799.
- Wang, L., Mao, M., Xie, J., Liao, Z., Zhang, H., and Li, H. (2023). Accurate solar pv power prediction interval method based on frequency-domain decomposition and lstm model. *Energy*, 262:125592.
- Wang, M., Peng, J., Luo, Y., Shen, Z., and Yang, H. (2021). Comparison of different simplistic prediction models for forecasting pv power output: assessment with experimental measurements. *Energy*, 224:120162.